arXiv:math/0312290v2 [math.RT] 10 Nov 2005

DESCRIPTION OF B–ORBIT CLOSURES OF ORDER 2 IN UPPER-TRIANGULAR MATRICES

ANNA MELNIKOV

Abstract. Let $\mathfrak{n}_n(\mathbb{C})$ be the algebra of strictly upper-triangular $n \times n$ matrices and $\mathcal{X}_2 = \{u \in \mathfrak{n}_n(\mathbb{C}) \ : \ u^2 = 0\}$ the subset of matrices of nilpotent order 2. Let $\mathbf{B}_n(\mathbb{C})$ be the group of invertible upper-triangular matrices acting on $\mathfrak{n}_n$ by conjugation. Let $\mathcal{B}_u$ be the orbit of $u \in \mathcal{X}_2$ with respect to this action. Let $\mathbf{S}_n^2$ be the subset of involutions in the symmetric group $\mathbf{S}_n$. We define a new partial order on $\mathbf{S}_n^2$ which gives the combinatorial description of the closure of $\mathcal{B}_u$. We also construct an ideal $\mathcal{I}(\mathcal{B}_u) \subset S(\mathfrak{n}^*)$ whose variety $\mathcal{V}(\mathcal{I}(\mathcal{B}_u))$ equals $\overline{\mathcal{B}}_u$.

We apply these results to orbital varieties of nilpotent order 2 in $\mathfrak{sl}_n(\mathbb{C})$ in order to give a complete combinatorial description of the closure of such an orbital variety in terms of Young tableaux. We also construct the ideal of definition of such an orbital variety up to taking the radical.

1. **Introduction**

1.1. Let $\mathbf{G} = \mathbf{SL}_n(\mathbb{C})$ be the special linear group of degree n over field $\mathbb{C}$. Let $\mathbf{B} = \mathbf{B}_n(\mathbb{C})$ be the standard Borel subgroup of $\mathbf{G}$, that is the subgroup of all invertible upper-triangular matrices.

Take $\mathfrak{g} = \mathfrak{sl}_n$ and let $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{h} \oplus \mathfrak{n}^-$ be its standard triangular decomposition where $\mathfrak{n} = \mathfrak{n}_n$ is a subalgebra of strictly upper-triangular matrices, $\mathfrak{h} = \mathfrak{h}_n$ is a subalgebra of diagonal matrices of trace zero and $\mathfrak{n}^- = \mathfrak{n}_n^-$ is a subalgebra of strictly lower-triangular matrices in $\mathfrak{sl}_n$.

We identify $\mathfrak{n}^-$ with $(\mathfrak{n}^+)^*$ through the Killing form on $\mathfrak{g}$. For any Lie algebra $\mathfrak{a}$, let $S(\mathfrak{a})$ denote its symmetric algebra.

For any $u \in \mathfrak{n}$ let $\mathcal{O}_u := \{AuA^{-1} \mid A \in \mathbf{G}\}$ be its (adjoint) $\mathbf{G}$-orbit and $\mathcal{B}_u := \{BuB^{-1} \mid B \in \mathbf{B}\}$ to be its (adjoint) $\mathbf{B}$-orbit.

For any $u \in \mathfrak{n}$, its only characteristic value is 0. So, $\mathcal{O}_u$ is a nilpotent orbit and every nilpotent orbit is obtained in such a way. Also, the Jordan form of a nilpotent orbit is described by a partition of n. Obviously, $\mathcal{B}_u \subset \mathcal{O}_u \cap \mathfrak{n}$. Let $\mathcal{O}$ be a nilpotent orbit and $v \in \mathfrak{n}$. We say that $\mathcal{B}_v$ is associated to $\mathcal{O}$ if $\mathcal{B}_v \subset \mathcal{O}$. It is obvious that $\mathcal{B}_v$ is associated to $\mathcal{O}$ if and only if $\mathcal{O}_v = \mathcal{O}$. As shown in [2] for $n \geq 6$, the number of $\mathcal{B}_v$ orbits associated

---

Supported in part by the Minerva Foundation, Germany, Grant No. 8466.

to $\mathcal{O}$ can be infinite. However, the $\mathbf{G}$ orbits of nilpotent order 2 decompose into finitely many $\mathbf{B}$ orbits and so are a nice exception.

Let $\mathcal{X}_2$ be the set of elements of nilpotent order 2 in $\mathfrak{n}$, that is $\mathcal{X}_2 := \{u \in \mathfrak{n} \mid u^2 = 0\}$ where u^2 is defined by associative multiplication of matrices. Let $\mathbf{S}_n$ be the symmetric group on n elements. Let $\mathbf{S}_n^2$ be the set of involutions in $\mathbf{S}_n$, that is $\mathbf{S}_n^2 := \{\sigma \in \mathbf{S}_n \mid \sigma^2 = Id\}$. As shown in [12] there is a natural bijection between $\mathbf{B}$-orbits in $\mathcal{X}_2$ and $\mathbf{S}_n^2$.

In this paper we define a new partial order on $\mathbf{S}_n^2$ which completely describes a $\mathbf{B}$-orbit closure for $\mathbf{B}$-orbits in $\mathcal{X}_2$ via this bijection. The combinatorial description of this order is natural and very simple. Moreover, for a given $\mathbf{B}$-orbit $\mathcal{B} \subset \mathcal{X}_2$ we construct (via generators) an ideal $\mathcal{I}(\mathcal{B}) \in S(\mathfrak{n}^-)$ whose variety $\mathcal{V}(\mathcal{I}(\mathcal{B}))$ equals $\overline{\mathcal{B}}$ and use it to prove our main Theorem 3.5.

In addition we apply these results to the study of orbital varieties in $\mathcal{X}_2$.

On the one hand, the ideas of N. Spaltenstein gave me the inspiration for the order defined here. On the other hand, we apply the results on $\mathbf{B}$-orbits to the study of orbital varieties. So we would like to give a brief description of orbital varieties.

1.2. Orbital varieties derive from the works of N. Spaltenstein [17, 18], and R. Steinberg [19, 20] written during their studies of the unipotent variety of a complex semisimple group $\mathbf{G}$. Translation from the unipotent variety of $\mathbf{G}$ to the nilpotent cone of $\mathfrak{g} = \mathrm{Lie}(\mathbf{G})$, gave the notion of an orbital variety. It is defined as follows.

Consider the adjoint action of $\mathbf{G}$ on $\mathfrak{g}$. Fix some triangular decomposition $\mathfrak{g} = \mathfrak{n} \bigoplus \mathfrak{h} \bigoplus \mathfrak{n}^-$. A $\mathbf{G}$ orbit $\mathcal{O}$ in $\mathfrak{g}$ is called nilpotent if it consists of nilpotent elements, that is if $\mathcal{O} = \mathbf{G}\,x$ for some $x \in \mathfrak{n}$. The intersection $\mathcal{O} \cap \mathfrak{n}$ is a Lagrangian subvariety of $\mathcal{O}$. It is reducible in general. Its irreducible components are called orbital varieties associated to $\mathcal{O}$. Orbital varieties play a key role in the study of primitive ideals in the enveloping algebra $U(\mathfrak{g})$. They also play an important role in Springer's Weyl group representations.

The first role above can be detailed as follows. Since $\mathfrak{g}$ is semisimple, we can identify $\mathfrak{g}^*$ with $\mathfrak{g}$ through the Killing form. This identification gives an adjoint orbit a symplectic structure. Then, by [18], [19] and [6], an orbital variety $\mathcal{U}$ associated to a nilpotent orbit is a Lagrangian subvariety. Following the orbit method one would like to attach an irreducible representation of the enveloping algebra $U(\mathfrak{g})$ to $\mathcal{U}$. This should be a simple highest weight module. Combining the results of A. Joseph and T. A. Springer one obtains a one to one correspondence between the set of primitive ideals of $U(\mathfrak{g})$ containing the augmentation ideal of its centre (thus corresponding to integral weights) and the set of orbital varieties in $\mathfrak{g}$ corresponding to Lusztig's special orbits (see for example [1]). The picture is especially beautiful for $\mathfrak{g} = \mathfrak{sl}_n$. In this case, all orbits are special and by [10] the associated variety of a simple highest (integral) weight module is irreducible. By [1] and [6] in general orbital variety closures are the irreducible components of an associated variety of a simple highest weight module. Whilst for $\mathfrak{g} = \mathfrak{sl}_n$ orbital variety closures are themselves associated varieties and therefore give a natural geometric understanding of the classification of primitive ideals. This makes their study especially interesting.

1.3. There are two descriptions of orbital varieties. The first one, valid only for $\mathfrak{sl}_n$ was given by Spaltenstein. The second one is a general construction for any semisimple Lie algebra. It was obtained by Steinberg. We will consider the first one in what follows. Both descriptions give a very nice combinatorial characterization of orbital varieties in $\mathfrak{sl}_n$ in terms of Young tableaux. However, both descriptions are not very satisfactory from a geometric point of view.

The ultimate description of an orbital variety $\mathcal{U}$ would be via the ideal $I(\mathcal{U}) \subset S(\mathfrak{n}^-)$ of definition of $\overline{\mathcal{U}}$, that is a radical ideal I whose variety $\mathcal{V}(I)$ equals $\overline{\mathcal{U}}$. In [7, Lecture 7] the general form that such an ideal should take is suggested. Let N be the generic matrix of $\mathfrak{n}^- + 1d$. Then $I(\mathcal{U})$ should be generated by $a_1, a_2, \cdots, a_k$. Here $a_i = \mathrm{gr}\,(m_i)$, where m_i are some minors of N and filtration is by degree. This suggestion derived from an algorithm for $\mathrm{Ann}_{\,U(\mathfrak{n}^-)}\mathcal{U}$ based on the Enright functor [6, 8.4] together with calculations in [2].

Of course, the above does not tackle the difficult question as to which minors m_i to choose. However, in [9] the ideal of definition of orbital varieties of codimension 1 in a nilradical was constructed. This is the simplest non-trivial case since such a variety closure is a complete intersection, thus we had to find only one non-trivial generating polynomial.

As explained in 1.1, here we apply the results for $\mathbf{B}$ orbits in $\mathcal{X}_2$, to orbital varieties associated to nilpotent orbits of nilpotent order 2. We use the fact shown in [11, 4.13] that an orbital variety $\mathcal{U} \subset \mathcal{X}_2$ admits a dense $\mathbf{B}$-orbit $\mathcal{B}_{\mathcal{U}}$. Applying our results to $\mathcal{B}_{\mathcal{U}}$ we obtain an ideal $\mathcal{I}(\mathcal{U}) := \mathcal{I}(\mathcal{B}_{\mathcal{U}})$ whose variety $\mathcal{V}(\mathcal{I}(\mathcal{U}))$ equals $\overline{\mathcal{U}}$. Then the ideal of definition of $\mathcal{U}$ is the radical of $\mathcal{I}(\mathcal{U})$.

1.4. Another natural problem connected to studying orbital variety closures is their description as a union of irreducible varieties each one lying in some nilpotent orbit. In particular, one can ask whether an orbital variety closure consists of orbital varieties only or it includes some other $\mathbf{B}$ stable varieties.

If $\mathfrak{g} \neq \mathfrak{sl}_n$ an orbital variety closure is not necessarily [15] a union of orbital varieties. However, the construction only works outside $\mathfrak{sl}_n$. One can conjecture that in $\mathfrak{sl}_n$ an orbital variety closure is a union of orbital varieties. This conjecture holds for orbital varieties of Richardson type as it is shown in [13]. As well it is supported by the computations for all orbital varieties for $n \leq 6$.

In this paper we show in 4.5 that this conjecture holds for orbital varieties in $\mathcal{X}_2$.

Given orbital varieties $\mathcal{U}, \mathcal{W} \in \mathfrak{n}$ we say that $\mathcal{W} \overset{\mathrm{G}}{\geq} \mathcal{U}$ if $\mathcal{W} \subset \overline{\mathcal{U}}$. In [14] we give the combinatorial description of this order for orbital varieties in $\mathcal{X}_2$ in terms of Young tableaux. Thus, the result in 4.5 completes the combinatorial description of an orbital variety closure in $\mathcal{X}_2$ in terms of Young tableaux. Although an orbital variety in $\mathcal{X}_2$ has a much simpler structure then an orbital variety in general even this relatively simple case already demonstrates the complexity of the theory.

1.5. The ideas we use to study $\overline{\mathcal{B}}_u$ are derived from Spaltenstein's characterization of orbital varieties in $\mathfrak{sl}_n$ by Young tableaux. Let us recall it in short.

A nilpotent orbit $\mathcal{O}_u$ is described by a partition giving the length of blocks in the Jordan form of u. Suppose the Jordan form of u consists of k non-zero blocks. We order blocks in non-increasing order and put λ_i^* to be the length of $i-$th block. Then $\mathcal{O}_u$ is defined by the partition $\lambda^* = (\lambda_1^*, \ldots, \lambda_k^*)$. We can represent λ^* graphically as a Young diagram – an array of k rows each starting at the same place on the left, where the i-th row contains λ_i^* boxes.

Since in this paper we consider only nilpotent elements with the Jordan blocks of length not more than 2, it is more convenient for us to take the dual partition to λ^* that is $\lambda = (\lambda_1, \ldots, \lambda_m)$ where $\lambda_i = \#\{\lambda_j^* \in \lambda^* \mid \lambda_j \geq i\}$. In these terms $\mathcal{O}_u$ is described by Young diagram D_λ with m columns with j-th column containing λ_j boxes. We will denote the corresponding orbit by $\mathcal{O}_\lambda$.

Recall that a Young tableau T associated to the Young diagram D_λ is obtained by filling the boxes of D_λ with numbers $1, \ldots, n$ so that the numbers increase in rows from left to right and in columns from top to bottom. Given a Young tableau T associated to D_λ its shape is defined to be λ and denoted as $\mathrm{sh}\,(T)$. Given $u \in \mathfrak{n} \cap \mathcal{O}_\lambda$ its Young diagram is again defined to be λ and denoted as $D_n(u)$, or simply $D(u)$. Now consider canonical projections $\pi_{1,n-i} : \mathfrak{n}_n \to \mathfrak{n}_{n-i}$ acting on a matrix by deleting the last i columns and the last i rows. Set $D_{n-1}(u) := D(\pi_{1,n-1}(u)), D_{n-2}(u) := D(\pi_{1,n-2}(u)), \ldots, D_1(u) := D(\pi_{1,1}(u))$ and $\phi(u) := (D_1(u), D_2(u), \ldots, D_n(u))$. Put 1 into the unique box of $D_1(u)$. For any $i\ : 1 \leq i \leq n-1$ note that $D_{i+1}(u)$ differs from $D_i(u)$ by a single "corner" box. Put $i+1$ into this box. This gives a bijection from the set of the chains $(D_1(u), D_2(u), \ldots, D_n(u))$ to the set of standard Young tableaux T of shape $D_n(u)$. In other words, we view a standard Young tableau as a chain of Young diagrams. For $u \in \mathfrak{n}$ put $\varphi(u) := T$ if T corresponds under this bijection to $\phi(u)$. Set $\nu_{_T} := \{u \in \mathfrak{n} \mid \varphi(u) = T\}$.

By Spaltenstein [17] orbital varieties associated to $\mathcal{O}_\lambda$ are parameterized by standard Young tableaux of shape λ as follows. Let $\{T_i\}$ be the set of Young tableaux of shape λ. Set $\mathcal{U}_{T_i} := \overline{\nu}_{_{T_i}} \cap \mathcal{O}_\lambda$. Then $\{\mathcal{U}_{T_i}\}$ is the set of orbital varieties associated to $\mathcal{O}_\lambda$.

1.6. We would like to push Spaltenstein's construction a little bit further. In a similar way define projections $\pi_{i,n} : \mathfrak{n} \to \mathfrak{n}_{\langle i,n\rangle}$ (cf. 2.3) acting on a matrix by deleting the first $i-1$ rows and the first $i-1$ columns. Set $D^i(u) := D(\pi_{n-i+1,n}(u))$ and $\theta(u) := (D^1(u), \ldots, D^n(u))$. Put n into the unique box of $D^1(u)$. Again, $D^{i+1}(u)$ differs from $D^i(u)$ by a single "corner" box. Insert $n-i+1$ into the left hand corner by inverting "jeu de taquin" (cf. [16]). This gives a second bijection from the set of chains $(D^1(u), D^2(u), \ldots, D^n(u))$ to the set of standard Young tableaux T of shape $D^n(u)$. In other words, we again view a standard Young tableau as a chain of Young diagrams. For $u \in \mathfrak{n}$ put $\vartheta(u) := T$ if T corresponds under this bijection to $\theta(u)$. Again, set $\nu^T := \{u \in \mathfrak{n} \mid \vartheta(u) = T\}$. By symmetry about the anti-diagonal it follows that any orbital variety $\mathcal{U}$ associated to $\mathcal{O}_\lambda$ is obtained as $\mathcal{U} = \overline{\nu^T} \cap \mathcal{O}_\lambda$. Moreover, it is immediate that $\overline{\nu^T} \cap \mathcal{O}_\lambda = \mathcal{U}_T$.

On the other hand, $\vartheta(u)$ can differ from $\varphi(u)$. For standard tableaux T, S of shape λ put $\nu_T^S = (\nu_T \cap \nu^S)$. Then $\nu_T = \coprod\limits_{\mathrm{sh}\, S=\lambda} (\nu_T \cap \nu^S)$. Obviously, $\nu_T^S \subset \mathcal{U}_T \cap \mathcal{U}_S$. Thus, $\dim \nu_T^S < \dim \mathcal{U}_T$ whenever $S \neq T$. Hence, ν_T^T is dense in $\mathcal{U}_T$.

For any $i, j : \ 1 \le i \le j \le n$ using jeu de taquin to delete $1, \ldots, i-1$ from a tableau T and deleting boxes containing $n, n-1, \ldots, j+1$ from T we define $\pi_{i,j}(T)$ – a tableau with $j-i+1$ boxes filled in with $i, \ldots, j$. Set $D_{i,j}(T) := \mathrm{sh}\,(\pi_{i,j}(T))$. In that way we can consider each Young tableau as a double chain through

$$T \longleftrightarrow \begin{bmatrix} D_{1,1}(T) & D_{1,2}(T) & \ldots & D_{1,n}(T) \\ & D_{2,2}(T) & \ldots & D_{2,n}(T) \\ & & \ddots & \vdots \\ & & & D_{n,n}(T) \end{bmatrix}$$

Correspondingly, we can define the projection $\pi_{i,j} : \mathfrak{n}_n \to \mathfrak{n}_{j-i+1}$ acting on a matrix by deleting the first $i-1$ rows and the first $i-1$ columns and the last $n-j$ rows and the last $n-j$ columns. Set

$$\nu'_T = \{u \in \mathfrak{n} \mid D(\pi_{i,j}(u)) = D_{i,j}(T)\}$$

Obviously, $\nu'_T \subset \nu_T^T$. Moreover, using Steinberg construction one gets that $\mathcal{U}_T = \overline{\nu'_T} \cap \mathcal{O}_\lambda$. This construction provides us the power rank conditions introduced in [21]. In turn these conditions define an ideal $\mathcal{I}_T \subset S(\mathfrak{n}^-)$ such that its variety $\mathcal{V}(\mathcal{I}_T)$ contains $\overline{\mathcal{U}}_T$. However, in general for $n \ge 6$ this ideal is far away from being an ideal of definition of $\mathcal{U}_T$ since the variety of the ideal is not irreducible and can include even another orbital variety associated to the same nilpotent orbit (cf., for example, [21]).

1.7. Now we can explain the way we construct $\mathcal{I}(\mathcal{B})$ for $\mathcal{B} \subset \mathcal{X}_2$. Note that adjoint $\mathbf{B}$ action is compatible projection, that is $\pi_{i,j}(\overline{\mathcal{B}}_u) = \overline{\mathcal{B}}'_{\pi_{i,j}(u)}$ where $\mathcal{B}'_{u'}$ is a $\mathbf{B}_{j-i+1}$–orbit of $u' \in \mathfrak{n}_{j-i+1}$. Thus, we can use the same construction of double chains not only for orbital varieties but for **B**-orbits as well. Note that a double chain defined by some Young tableau corresponds in general to some set of **B**-orbits, however, a double chain defined by a **B**-orbit will not correspond in general to a standard Young tableau. This double chain can be described by power rank conditions exactly in the same way as in the case of an orbital variety. Thus, as in the case of orbital varieties we can construct an ideal $\mathcal{I}(\mathcal{B}) \subset S(\mathfrak{n}^-)$ as it is explained in detail in 2.8, 2.9. Again, the variety $\mathcal{V}(\mathcal{I}(\mathcal{B}))$ always contains $\overline{\mathcal{B}}$ but in general it contains $\mathbf{B}$–orbits not from $\overline{\mathcal{B}}$ as well (cf. 2.10).

However, for $\mathcal{B} \subset \mathcal{X}^2$ the variety $\mathcal{V}(\mathcal{I}(\mathcal{B})) = \overline{\mathcal{B}}$. This is the main technical result of the paper.

1.8. In this paper we do not touch upon two very important problems connected to our results. The first problem is whether the ideals we construct are ideals of definition. To show this we have to show that they are radical ideals. This is a complex question. For some cases we can answer positively using technique introduced in [4], in particular if $\mathcal{B}$ is associated to the minimal (non-zero) orbit. Indeed, for the nilpotent orbit of the smallest non-zero dimension, the ideal constructed in 2.9 for an orbital variety associated to it, coincides with the ideal of definiton for such orbital varieties given in [3]. These computations give us a hope that our ideals are ideals of definition. However, I am far away from complete answer to this question.

The second problem is closely connected to the first one. It is the construction of a strong quantization of orbital varieties in $\mathcal{X}_2$. Again, as it is shown in [8] the problem is far from being easy even for the minimal nilpotent orbit.

Another question, that we do not consider here, which however could be solved using our construction is the description of intersections of orbital varieties of nilpotent order 2.

1.9. The body of the paper consists of three sections. In section 2 we explain all the background essential in the subsequent analysis to make the paper self-contained. In section 3 we formulate and prove the results on **B**-orbit closures in $\mathcal{X}_2$. Finally, in section 4 we explain the results connected to orbital varieties of nilpotent order 2 and apply the results of section 3 to complete the combinatorial description of their closures.

In the end one can find the index of notation in which symbols appearing frequently are given with the subsection where they are defined. We hope that this will help the reader to find his way through the paper.

Acknowledgments. This research was done during my stay as a guest at the Weizmann Institute. I am very grateful to my host A. Joseph for the invitation and many fruitful discussions through the various stages of this work.

I would also like to express my gratitude to the referee. His numerous remarks helped to improve the presentation and alter some proofs. His remarks and the remarks of A. Joseph helped me to bring this paper to a more clear and, hopefully, readable form.

2. **Geometric and Combinatorial Preliminaries**

2.1. Let $P(n)$ denote the set of partitions of n. For all $\lambda := \{\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_k > 0\} \in P(n)$ we define the Young diagram D_λ to be the array of k **columns** of boxes each starting at the first row with the i-th column containing λ_i boxes. Let $\mathbf{D}_n$ denote the set of all Young diagrams with n boxes.

Recall as well that λ^* denotes the dual partition.

For example, take $\lambda = (4,2,2)$. Then $\lambda^* = (3,3,1,1)$ and

$D_\lambda =$



2.2. It is convenient to replace $\mathfrak{sl}_n$ by $\mathfrak{gl}_n$. This makes no difference to the nilpotent cone $\mathcal{N}$ whilst the adjoint action (conjugation) by $\mathbf{G} = \mathbf{GL}_n$ on $\mathcal{N}$ factors to $\mathbf{SL}_n$. Let $\mathbf{B}$ be the (Borel) subgroup of upper-triangular matrices in $\mathbf{G}$. Let $\mathfrak{b} = \operatorname{Lie}(\mathbf{B}) = \mathfrak{n} \oplus \mathfrak{h}$ be the corresponding subalgebra of $\mathfrak{gl}_n$.

Recall the $\mathcal{O}_u$ defined in 1.1 and the $\mathcal{O}_\lambda$ defined in 1.5. For $u \in \mathfrak{n} \cap \mathcal{O}_\lambda$ set $\operatorname{sh}(u) := \lambda$ and $\operatorname{sh}(\mathcal{O}_u) := \operatorname{sh}(u)$.

In particular, $\mathcal{O}_{(l,k)}$ denotes the nilpotent orbit with k Jordan blocks of length 2 and $l-k$ Jordan blocks of length 1.

2.3. Let $e_{i,j}$ be a matrix having 1 in the ij-th entry and 0 elsewhere. Then $\{e_{i,j}\}_{i,j=1}^n$ is a basis of $\mathfrak{g}$.

Let $R \subset \mathfrak{h}^*$ denote the set of non-zero roots, R^+ the set of positive roots corresponding to $\mathfrak{n}$ and $\Pi \subset R^+$ the resulting set of simple roots.

Take $i < j$ and let $\alpha_{i,j}$ be the root which is the weight of $e_{i,j}$. Set $\alpha_{j,i} = -\alpha_{i,j}$. We write $\alpha_{i,i+1}$ simply as α_i. Then $\Pi = \{\alpha_i\}_{i=1}^{n-1}$. Moreover, $\alpha_{i,j} \in R^+ \iff i < j$. One has

$$\alpha_{i,j} = \begin{cases} \sum\limits_{k=i}^{j-1} \alpha_k & \text{if } i > j \\ -\sum\limits_{k=i}^{j-1} \alpha_k & \text{if } i < j \end{cases}$$

Let $X_{\alpha_{i,j}} := X_{i,j} := \mathbb{C}e_{i,j}$ be the root space defined by $\alpha_{i,j} \in R$.

For $\alpha_i \in \Pi$, let $\mathbf{P}_{\alpha_i}$ be the standard parabolic subgroup of $\mathbf{G}$ with $\operatorname{Lie}(P_{\alpha_i}) = \mathfrak{b} \oplus X_{-\alpha_i} = \mathfrak{b} \oplus X_{i+1,i}$.

For $1 \leq i < j \leq n$ let $\mathbf{P}_{\langle i,j \rangle}$ be the standard parabolic subgroup of $\mathbf{G}$ such that $\mathbf{B}, \{\mathbf{P}_{\alpha_k}\}_{k=i}^{j-1} \subset \mathbf{P}_{\langle i,j \rangle}$, equivalently such that

$$\operatorname{Lie}(\mathbf{P}_{\langle i,j \rangle}) = \mathfrak{b} \oplus \bigoplus_{i \leq s < t \leq j} X_{t,s}$$

Let $\mathbf{M}_{\langle i,j \rangle}$ be the unipotent radical of $\mathbf{P}_{\langle i,j \rangle}$ and $\mathbf{L}_{\langle i,j \rangle}$ its Levi factor. Set $\mathbf{B}_{\langle i,j \rangle} := \mathbf{B} \cap \mathbf{L}_{\langle i,j \rangle}$. Set $\mathfrak{m}_{\langle i,j \rangle} := \operatorname{Lie}(\mathbf{M}_{\langle i,j \rangle})$ and $\mathfrak{l}_{\langle i,j \rangle} := \operatorname{Lie}(\mathbf{L}_{\langle i,j \rangle})$. Set $\mathfrak{n}_{\langle i,j \rangle} := \bigoplus\limits_{i \leq s < t \leq j} X_{k,s}$ (that is $\mathfrak{n}_{\langle i,j \rangle} = \mathfrak{n} \cap \mathfrak{l}_{\langle i,j \rangle})$). We have decompositions $\mathbf{B} = \mathbf{M}_{\langle i,j \rangle} \ltimes \mathbf{B}_{\langle i,j \rangle}$ and $\mathfrak{n} = \mathfrak{n}_{\langle i,j \rangle} \oplus \mathfrak{m}_{\langle i,j \rangle}$. They define projections $\pi_{i,j} : \mathbf{B} \to \mathbf{B}_{\langle i,j\rangle}$ and $\pi_{i,j} : \mathfrak{n} \to \mathfrak{n}_{\langle i,j\rangle}$ which we mentioned in 1.6. For the matrix $u = (u_{s,t})_{s,t=1}^n$ where $u \in \mathfrak{n}$ or $u \in \mathbf{B}$ one has

$$\pi_{i,j}(u) = \begin{pmatrix} u_{i,i} & u_{i,i+1} & \dots & u_{i,j} \\ 0 & u_{i+1,i+1} & \dots & u_{i+1,j} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & u_{j,j} \end{pmatrix}$$

Recall 1.6. For $u \in \mathfrak{n}$ put $D_{i,j}(u) := \mathrm{sh}\,(\pi_{i,j}(u))$. Let D_u be the resulting "matrix" of projections shapes, that is

$$(D_u)_{i,j} = \begin{cases} 0, & \text{if } i \geq j; \\ D_{i,j}(u) & \text{otherwise} \end{cases}$$

Let $\mathcal{B}_u$ denote the $\mathbf{B}$ orbit of u.

For any $u \in \mathfrak{n}$ and $A \in \mathbf{B}$ one has $\pi_{i,j}(AuA^{-1}) = \pi_{i,j}(A)\pi_{i,j}(u)\pi_{i,j}(A)^{-1}$ so that $\mathrm{sh}\,(\pi_{i,j}(AuA^{-1})) = \mathrm{sh}\,(\pi_{i,j}(u))$ and $D_{\mathcal{B}_u} := D_u$ is well defined. By the discussion in 1.6 one sees that $D_{i,j-1}(u)$ (as well as $D_{i+1,j}$) is obtained from $D_{i,j}(u)$ just by decreasing one of the entries of $D_{i,j}$ by 1, that is if $D_{i,j}(u) = (\lambda_1, \dots, \lambda_k)$, where $\lambda_k > 0$, then there exists $r :\ 1 \leq r \leq k$ such that $D_{i,j-1}(u) = (\lambda_1, \dots, \lambda_r - 1, \dots, \lambda_k)$.

For example,

$$u = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \quad D_{\mathcal{B}_u} = D_u = \begin{pmatrix} 0 & (1,1) & (2,1) & (2,1,1) & (2,2,1) \\ 0 & 0 & (2) & (2,1) & (2,2) \\ 0 & 0 & 0 & (2) & (2,1) \\ 0 & 0 & 0 & 0 & (2) \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

2.4. Let W be Weyl group of $(\mathfrak{g}, \mathfrak{h})$. For $\alpha \in R^+$ let $s_\alpha \in W$ be the corresponding reflection. For $\alpha_i \in \Pi$ set $s_i := s_{\alpha_i}$. We identify W with $\mathbf{S}_n$ by taking s_i to be the elementary permutation interchanging i, $i+1$.

Set $\mathbf{S}_{\langle i,j\rangle} := \langle s_{\alpha_k} \ :\ i \leq k < j \ \rangle$ which we call a parabolic subgroup of $W = \mathbf{S}_n$. Set $D_{\langle i,j\rangle} := \{w \in \mathbf{S}_n \mid w(\alpha_k) \in R^+ \ \forall i \leq k < j\}$. Then the multiplication map gives a bijection $\mathbf{S}_{\langle i,j\rangle} \times D_{\langle i,j\rangle} \simeq \mathbf{S}_n$. Let $\pi_{i,j} : \mathbf{S}_n \to \mathbf{S}_{\langle i,j\rangle}$ be the projection onto the first factor.

2.5. Let us define the decomposition of $\mathcal{X}_2$ into $\mathbf{B}-$orbits.

Recall that $\mathbf{S}_n^2$ denotes the subset of involutions in $\mathbf{S}_n$. Write $\sigma \in \mathbf{S}_n^2$ as a product of disjoint cycles of length 2. Order entries inside a given cycle to be increasing. Unless mentioned to the contrary (in section 4) we order the cycles so that the first entries increase. Thus, $\sigma = (i_1, j_1)(i_2, j_2) \dots (i_k, j_k)$ where $i_s < j_s$ for any $1 \leq s \leq k$ and $i_s < i_{s+1}$ for any $1 \leq s < k$.

For $\sigma=(i_1,j_1)(i_2,j_2)\ldots(i_k,j_k)$ we construct $N_\sigma\in\mathfrak{n}$ as follows

$$(N_\sigma)_{i,j}=\begin{cases}1, & \text{if } i=i_s, j=j_s \text{ for some } 1\le s\le k:\\ 0, & \text{otherwise.}\end{cases}$$

Note that N_σ is the upper-triangular part of the permutation matrix M_σ. Set $\mathcal{B}_\sigma:=\mathcal{B}_{N_\sigma}$. By [12, 2.2] one has

Proposition. $\mathcal{X}_2=\coprod\limits_{\sigma\in\mathbf{S}_n^2}\mathcal{B}_\sigma$.

This finiteness property is particular for $\mathcal{X}_2$. For $n\ge 6$ and $\mathcal{X}_k=\{X\in\mathfrak{n}\mid X^k=0\}$ where $k\ge 3$ the number of **B** orbits in $\mathcal{X}_k$ is infinite in general for any infinite field as it is shown in [2].

2.6. Given $\sigma=(i_1,j_1)(i_2,j_2)\ldots(i_k,j_k)\in\mathbf{S}_n^2$, set $l(\sigma):=k$ and let $\mathcal{O}_\sigma:=\mathcal{O}_{N_\sigma}$ be the **G** orbit of N_σ.

Lemma. *For $\sigma\in\mathbf{S}_n^2$ such that $l(\sigma)=k$ one has $\mathrm{sh}\,(\mathcal{O}_\sigma)=(n-k,k)$.*

Proof.

Indeed, $N_\sigma\in\mathcal{X}_2$ and its rank is k. Thus, its Jordan form contains k blocks of length 2 and $n-k$ blocks of length 1.

■

2.7. Given $\sigma=(i_1,j_1)(i_2,j_2)\ldots(i_k,j_k)$ ordered as in 2.5. For $s\ :2\le s\le k$ set

$$r_s:=r(i_s,j_s):=\#\{j_p\ :\ p<s,\ j_p<j_s\}+\#\{j_p\ :\ j_p<i_s\}.$$

For example, take $\sigma=(1,6)(3,4)(5,7)$. Then $l(\sigma)=3$ and $r_2=0,\ r_3=2+1=3$.

By [12, 3.1] one has

Theorem. *For $\sigma=(i_1,j_1)(i_2,j_2)\ldots(i_k,j_k)\in\mathbf{S}_n^2$ one has*

$$\dim\mathcal{B}_\sigma=kn+\sum_{s=1}^{k}(i_s-j_s)-\sum_{s=2}^{k}r_s.$$

2.8. Given $\lambda=(\lambda_1,\ldots,\lambda_k)$ and $\mu=(\mu_1,\ldots,\mu_l)$ in $P(n)$ we define $\lambda\le\mu$ if for all $s\ :\ 1\le s\le\min(k,l)$ one has

$$\sum_{i=1}^{s}\lambda_i\le\sum_{i=1}^{s}\mu_i.$$

By a result of Gerstenhaber (cf. [5], for example) one has

$$\overline{\mathcal{O}_\lambda}=\coprod_{\mu\ge\lambda}\mathcal{O}_\mu.$$

By the Jordan form, $\mathrm{sh}\,(\mathcal{O}_u)=(\lambda_1,\dots,\lambda_k)$ if and only if

$$\mathrm{Rank}\,u=n-\lambda_1,\ \mathrm{Rank}\,u^2=n-(\lambda_1+\lambda_2),\dots,\mathrm{Rank}\,u^{k-1}=\lambda_k,\ \mathrm{Rank}\,u^k=0$$

Moreover, by the above result of Gerstenhaber $u'\in\overline{\mathcal{O}}_u$, if and only if

$$\mathrm{Rank}\,u'\leq n-\lambda_1=\sum_{s=2}^{k}\lambda_s,\ \mathrm{Rank}\,{u'}^2\leq n-(\lambda_1+\lambda_2)=\sum_{s=3}^{k}\lambda_s,\dots,$$

$$\mathrm{Rank}\,{u'}^{k-1}\leq\lambda_k,\ \mathrm{Rank}\,{u'}^k=0$$

We call these the power rank conditions. Each power rank condition can be translated into polynomials in $S(\mathfrak{g}^*)$ as follows. Let us identify $S(\mathfrak{g}^*)$ with $S(\mathfrak{g})$ through the coordinate functions

$$x_{j,i}(e_{r,s})=\begin{cases}1, & \text{if } (s,r)=(i,j),\\ 0, & \text{otherwise.}\end{cases}\tag{$*$}$$

Setting $x_{i,j}=e_{j,i}$ identifies $\mathfrak{g}^*$ with $\mathfrak{g}$. Now, $\mathrm{Rank}\,u^i\leq j$ if and only if every minor of order $j+1$ of u^i is equal to 0. Let N be the matrix with indeterminate $x_{i,j}$ in the ij-th place . Let $\{P_t^{(i,j)}\}_{t=1}^{\binom{n}{j}^2}$ be the set of all minors of order j of N^i. Then $\overline{\mathcal{O}}_\lambda$ is the zero variety of the ideal of $S(\mathfrak{g}^*)$ generated by $\left\{P_t^{(i,j_i)}\right\}_{i=1,\ t=1}^{k,\ \binom{n}{j_i}^2}$, where $j_i=1+\sum\limits_{s=i+1}^{k}\lambda_s$.

2.9. Identify $S(\mathfrak{n}^*)$ with $S(\mathfrak{n}^-)$ through 2.8 $(*)$. Take $u\in\mathfrak{n}$ and put $\mathcal{B}:=\mathcal{B}_u$. Let us apply 2.8 to $\mathcal{B}$.

Let $D_\mathcal{B}$ be the matrix defined in 2.3. From $D_\mathcal{B}$ define an ideal $I(\mathcal{B})$ (or simply I) of $S(\mathfrak{n}^-)$ as follows. Suppose $(D_\mathcal{B})_{i,j}=(\lambda_1(i,j),\dots,\lambda_k(i,j))$. For any $u\in\mathcal{B}$ one has by definition of $D_\mathcal{B}$ that the power rank conditions on $\pi_{i,j}(u)$ (cf. [21]) are as follows:

$$\mathrm{Rank}\,(\pi_{i,j}(u))=j-i+1-\lambda_1(i,j),\ \mathrm{Rank}\,((\pi_{i,j}(u))^2)=j-i+1-\lambda_1(i,j)-\lambda_2(i,j),\ \dots$$

$$\mathrm{Rank}\,((\pi_{i,j}(u))^{k-1})=\lambda_k(i,j),\quad \mathrm{Rank}\,((\pi_{i,j}(u))^k)=0.\tag{$*$}$$

In turn, these power rank conditions translate to polynomials in $S(\mathfrak{n}^-)$ by 2.8.

Note that if $u\in\mathcal{X}_2$ then the power rank conditions provide us polynomials of 2 types: quadratic polynomials given by the conditions $\pi_{i,j}(u^2)=0$ and polynomials of order $\mathrm{Rank}\,(\pi_{i,j}(u))+1$ – these are the minors of order $\mathrm{Rank}\,(\pi_{i,j}(u))+1$ of matrix $\pi_{i,j}(u)$ (cf. 3.4 for the details).

Note that for any $u\in\mathfrak{n}$ one has

(i) $\pi_{i,j}(u^r)=(\pi_{i,j}(u))^r$ for any $i,j\ :1\leq i<j\leq n$ and $r\in\mathbb{N}$.
(ii) $\mathrm{Rank}\,(\pi_{i,j}(u))\leq\mathrm{Rank}\,(\pi_{i,j-1}(u))+1$ and $\mathrm{Rank}\,(\pi_{i,j}(u))\leq\mathrm{Rank}\,(\pi_{i+1,j}(u))+1$ for any possible i,j.

Let $\mathcal{I}_{(i,j)}(\mathcal{B})$ be the ideal generated by power rank conditions of $(D_{\mathcal{B}})_{i,j}$.

For $r\ :\ 1 \le r \le n-1$ and $i\ :\ 1 \le i \le n-r$ let us call the set of elements $(i, i+r)$ of a matrix, its $r-$th set. Let $\mathcal{I}_r(\mathcal{B})$ be the ideal generated by the power rank conditions (translated into corresponding minors) of sets $1, \ldots, r$, of $D_{\mathcal{B}}$ (so that $\mathcal{I}_r(\mathcal{B}) = \mathcal{I}_{r-1}(\mathcal{B}) + \sum_{i=1}^{n-r} \mathcal{I}_{(i,i+r)}(\mathcal{B})$ and $\mathcal{I}(\mathcal{B}) = \mathcal{I}_{n-1}(\mathcal{B})$).

Lemma. *Let $\mathcal{B}$ be a $\mathbf{B}$ orbit in $\mathfrak{n}$. Let $\mathcal{I}_{r-1}(\mathcal{B})$ be the ideal generated by the power rank conditions of sets $1, \ldots, r-1$ of $D_{\mathcal{B}}$ Let $\lambda = (\lambda_1, \lambda_2, \ldots)$ be either the partition $D_{i,i+r-1}$ or $D_{i+1,i+r}$. For some $l \in \mathbb{N}$ we can write $D_{i,i+r} = (\lambda_1, \ldots, \lambda_l + 1, \ldots)$. Then the polynomials generated by power rank conditions coming from $D_{i,i+r}$ for powers less than l belong to the ideal $\mathcal{I}_{r-1}$. In particular,*

(i) *if $D_{i,i+r} = (\lambda_1, \ldots, \lambda_k, 1)$ then $\mathcal{I}_{(i,i+r)}(\mathcal{B}) \subset \mathcal{I}_{r-1}(\mathcal{B})$;*

(ii) *if $\lambda = (\lambda_1, \lambda_2)$ and $D_{i,i+r} = (\lambda_1, \lambda_2 + 1)$ (that is $l = 2$) then $\mathcal{I}_{r-1} + \mathcal{I}_{(i,i+r)} = \mathcal{I}_{r-1} + \langle \sum_{s=i+1}^{i+r-1} x_{i+r,s} x_{s,i} \rangle$*

Proof.

Suppose $D_{i,i+r-1} = (\lambda_1, \ldots, \lambda_k)$ then $D_{i,i+r} = (\mu_1, \ldots, \mu_k,)$ where $\mu_l = \lambda_l + 1$ for some $l \in \mathbb{N}$ and $\mu_s = \lambda_s$ for any $s \ne l$. The power rank conditions give us

$$\operatorname{Rank}\left((\pi_{i,i+r}(u))^p\right) = \sum_{t=p+1}^{k} \mu_t = \begin{cases} \sum_{t=p+1}^{k} \lambda_t + 1, & \text{if } p < l; \\ \sum_{t=p+1}^{k} \lambda_t, & \text{if } p \ge l; \end{cases}$$

Define N as in 2.8. Let $\{m_{i,j}^p(N^s)\}$ be the set of minors of order p of $(\pi_{i,j}(N))^s$, where $\pi_{i,j}(N)$ is obtained by deleting the first $i-1$ rows and columns and the last $n-j$ rows and columns. Since N is upper-triangular one has $\pi_{i,j}(N^s) = (\pi_{i,j}(N))^s$. Then the condition $\operatorname{Rank}\left((\pi_{i,i+r}(u)^p)\right) = \sum_{t=p+1}^{k} \mu_t$ gives the subset of minors $\{m_{i,i+r}^{1+\sum_{t=p+1}^{k} \mu_t}(N^p)\}$. If $p < l$ one has $m_{i,i+r-1}^{1+\sum_{t=p+1}^{k} \lambda_t}(N^p) \in \mathcal{I}_{r-1}(\mathcal{B})$. Taking the expansion of $m_{i,i+r}^{1+\sum_{t=p+1}^{k} \mu_t}(N^p)$ by its last column we get minors $\{m_{i,i+r-1}^{1+\sum_{t=p+1}^{k} \lambda_t}(N^p)\}$ in the sum so that $m_{i,i+r}^{1+\sum_{t=p+1}^{k} \mu_t}(N^p) \in \mathcal{I}_{r-1}(\mathcal{B})$.

■

Let us determine $\mathcal{I}(\mathcal{B})$ for the example in 2.3. Recall that

$$D_{\mathcal{B}_u} = \begin{pmatrix} 0 & (1,1) & (2,1) & (2,1,1) & (2,2,1) \\ 0 & 0 & (2) & (2,1) & (2,2) \\ 0 & 0 & 0 & (2) & (2,1) \\ 0 & 0 & 0 & 0 & (2) \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

In our notation we get $\mathcal{I}_1(\mathcal{B}) = \langle P_1 = x_{3,2},\ P_2 = x_{4,3}, P_3 = x_{5,4} \rangle$. As for the entries of the second set one has $D_{i,i+2} = (2,1) = (D_{i+1,i+2}, 1)$ for all $i\ :\ 1 \le i \le 3$, thus, the second set does not add new generating polynomials, so that $\mathcal{I}_2(\mathcal{B}) = \mathcal{I}_1(\mathcal{B})$. Again, since $D_{1,4} = (2,1,1) = (D_{1,3}, 1)$ then by our lemma we do not get new generating polynomials from this condition. The only possible new generating polynomial produced by the third set obtains from $D_{2,5} = (2,2)$ and it comes from $\mathrm{Rank}\,(\pi_{2,5}(N^2)) = 0$. This gives us $P_3 = x_{5,3}x_{3,2} + x_{5,4}x_{4,2}$ however $P_3 = P_1x_{5,3} + x_{4,2}P_3$ so that it is in $\mathcal{I}_1(\mathcal{B})$. Finally, $D_{1,5} = (2,2,1) = (D_{2,5},1)$ so that again by our lemma we do not get new generating polynomials. Thus, $\mathcal{I}(\mathcal{B}) = \mathcal{I}_1(\mathcal{B}) = \langle x_{3,2}, x_{4,3}, x_{5,4}\rangle$.

2.10. Given two $\mathbf{B}-$orbits $\mathcal{B}, \mathcal{B}'$ applying the result of Gerstenhaber we get $\mathcal{I}(\mathcal{B}) \subset \mathcal{I}(\mathcal{B}')$ if and only if for any $i < j$ one has $(D_{\mathcal{B}})_{i,j} \leq (D_{\mathcal{B}'})_{i,j}$.

Obviously, $\overline{\mathcal{B}} \subset \mathcal{V}(\mathcal{I}(\mathcal{B}))$. However, in general $\mathcal{V}(\mathcal{I}(\mathcal{B})) \supsetneq \overline{\mathcal{B}}$. Power rank conditions are too weak even to separate two $\mathbf{B}$-orbits of the same dimension lying in the same nilpotent orbit $\mathcal{O}$. For example, consider

$$M = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}, \qquad \text{and} \qquad N = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

One has $M, N \in \mathcal{O}_{(3,2,1)}$ and $\dim \mathcal{B}_N = \dim \mathcal{B}_M = 11$ thus $\overline{\mathcal{B}}_N \not\subset \overline{\mathcal{B}}_M$. On the other hand

$$D_M = \begin{pmatrix} 0 & (1,1) & (2,1) & (2,2) & (2,2,1) & (3,2,1) \\ 0 & 0 & (2) & (2,1) & (2,1,1) & (3,1,1) \\ 0 & 0 & 0 & (1,1) & (1,1,1) & (2,1,1) \\ 0 & 0 & 0 & 0 & (1,1) & (2,1) \\ 0 & 0 & 0 & 0 & 0 & (2) \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$D_N = \begin{pmatrix} 0 & (1,1) & (2,1) & (3,1) & (3,1,1) & (3,2,1) \\ 0 & 0 & (2) & (3) & (3,1) & (3,2) \\ 0 & 0 & 0 & (2) & (2,1) & (2,2) \\ 0 & 0 & 0 & 0 & (1,1) & (2,1) \\ 0 & 0 & 0 & 0 & 0 & (2) \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

So that $(D_M)_{i,j} \geq (D_N)_{i,j}$ for all possible i,j.Thus, $\mathcal{I}(\mathcal{B}_M) \subsetneq \mathcal{I}(\mathcal{B}_N)$ so that $\overline{\mathcal{B}}_N \subset \mathcal{V}(\mathcal{I}(\mathcal{B}_N)) \subset \mathcal{V}(\mathcal{I}(\mathcal{B}_M))$. Hence, $\mathcal{V}(\mathcal{I}(\mathcal{B}_M)) \supsetneq \overline{\mathcal{B}}_M$.

2.11. From the previous subsections we see that we can define two partial orders on $\mathbf{B}$ orbits in $\mathfrak{n}$. The first one is defined by inclusion of $\mathbf{B}$ orbit closures and the second one by inverse inclusion of the ideals generated by power rank conditions defined by a $\mathbf{B}$ orbit. We call the first order geometric and the second order algebraic. From 2.10 we see that in general the algebraic order is a proper extension of the geometric order.

Now consider just $\mathbf{B}$ orbits of nilpotent order 2 in $\mathfrak{n}$. They are in bijection with $\mathbf{S}_n^2$ so both these orders induce partial orders on $\mathbf{S}_n^2$.

The geometric order on $\mathbf{S}_n^2$ is induced by $\mathbf{B}$ orbit closures inclusion, that is for $\sigma, \sigma' \in \mathbf{S}_n^2$ we set $\sigma' \overset{\mathrm{G}}{\leq} \sigma$ if $\overline{\mathcal{B}}_{\sigma'} \subset \overline{\mathcal{B}}_{\sigma}$.

The algebraic order on $\mathbf{S}_n^2$ is induced by inverse inclusions of $\mathcal{I}(\mathcal{B}_\sigma)$, that is for $\sigma, \sigma' \in \mathbf{S}_n^2$ we set $\sigma' \overset{\mathrm{A}}{\leq} \sigma$ if $\mathcal{I}(\mathcal{B}_{\sigma'}) \supset \mathcal{I}(\mathcal{B}_\sigma)$ (or equivalently if $\mathcal{V}(\mathcal{I}(\mathcal{B}_{\sigma'})) \subset \mathcal{V}(\mathcal{I}(\mathcal{B}_\sigma))$).

We show in what follows that on $\mathbf{S}_n^2$ the geometric and the algebraic orders coincide and we have the precise combinatorial description of this order.

2.12. Since $\pi_{i,j}$ is a projection it is continuous, one has

Lemma. *For* $\mathbf{B}$*-orbit* $\mathcal{B}$ *and any* $i,j\ :\ 1 \leq i < j \leq n$ *one has* $\pi_{i,j}(\overline{\mathcal{B}}) \subseteq \overline{\pi_{i,j}(\mathcal{B})}$. *In particular, if* $\mathcal{B}' \subset \overline{\mathcal{B}}$ *then* $\pi_{i,j}(\mathcal{B}') \subset \overline{\pi_{i,j}(\mathcal{B})}$.

2.13. We need also the following notation for $\sigma \in \mathbf{S}_n^2$. We write $(i,j) \in \sigma$ if it is one of the cycles of σ.

We write $\sigma = (i_1, j_1) \ldots (i_k, j_k)$ in increasing order as described in 2.5. Set $I(\sigma) := \{i_s\}_{s=1}^k$ and $J(\sigma) := \{j_s\}_{s=1}^k$.

For $1 \leq i < j \leq n$ set $S_{i,j}$ to be the set of indices $S_{i,j}(\sigma) := \{s\ :\ i \leq i_s \leq j \text{ and } i \leq j_s \leq j\}$. For example, take $\sigma = (1,8)(2,5)(3,4)(6,7) \in \mathbf{S}_8$. Then $S_{2,6}(\sigma) = \{2,3\}$. One has $\pi_{i,j}(\sigma) = (i_{s_1}, j_{s_1}) \ldots (i_{s_q}, j_{s_q})$, where $S_{i,j} = \{s_p\}_{p=1}^q$. In our example $\pi_{2,6}(\sigma) = (2,5)(3,4)$.

3. B-orbit closures in $\mathcal{X}_2$

3.1. For a matrix $u \in \mathfrak{n}$ recall the notion $\mathcal{B}_u$ from 1.1 and $D_u, D_{\mathcal{B}_u}$ from 2.3. For $i,j\ :\ 1 \leq i < j \leq n$ set $R_{i,j}(u) = \operatorname{Rank} \pi_{i,j}(u)$ and let R_u denote the corresponding (upper-triangular) matrix of ranks:

$$(R_u)_{i,j} := \begin{cases} 0, & \text{if } i \geq j; \\ R_{i,j}(u), & \text{otherwise.} \end{cases}$$

Obviously, for any $v \in \mathcal{B}_u$ one has $R_v = R_u$ so that we may write $R_{\mathcal{B}_u} := R_u$ exactly as we have defined $D_{\mathcal{B}_u}$.

Note that R_u can be read off from D_u. Indeed, denote $D_{i,j}(u) := (\lambda^1_{i,j}(u), \lambda^2_{i,j}(u), \ldots)$ then $R_{i,j}(u) = j + 1 - i - \lambda^1_{i,j}(u)$. In general, the information encoded in D_u is much reacher than the information encoded in R_u. However, in the case of $u \in \mathcal{X}_2$ these two matrices give exactly the same information since $D_{i,j}(u) = (j + 1 - i - R_{i,j}(u),\ R_{i,j}(u))$.

Adding one row or one column to a matrix can at most increase its rank by one, so for any $u \in \mathfrak{n}$ the matrix R_u has the following properties

(i) $(R_u)_{i,j} = 0$ if $i \geq j$

(ii) For $i < j$ one has $(R_u)_{i+1,j} \leq (R_u)_{i,j} \leq (R_u)_{i+1,j} + 1$ and $(R_u)_{i,j-1} \leq (R_u)_{i,j} \leq (R_u)_{i,j-1} + 1$.

By 2.5 for any $u \in \mathcal{X}_2$ there exist a unique $\sigma \in \mathbf{S}_n^2$ such that $\mathcal{B}_u = \mathcal{B}_\sigma$. We get $R_u = R_{N_\sigma}$ which we denote by $R_\sigma := R_{N_\sigma} = R_{\mathcal{B}_\sigma}$. By the definition of N_σ in each row and each column there is not more than one non-zero entry, thus, $(R_\sigma)_{i,j}$ is simply the number of ones in the matrix $\pi_{i,j}(N_\sigma)$.

For example, take $\sigma = (1,7)(2,3)(4,6) \in \mathbf{S}_8$. Then

$$N_\sigma = \begin{pmatrix} 0&0&0&0&0&0&1&0\\ 0&0&1&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&1&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0 \end{pmatrix} \quad \text{and} \quad R_\sigma = \begin{pmatrix} 0&0&1&1&1&2&3&3\\ 0&0&1&1&1&2&2&2\\ 0&0&0&0&0&1&1&1\\ 0&0&0&0&0&1&1&1\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0\\ 0&0&0&0&0&0&0&0 \end{pmatrix}$$

In particular, just by the form of N_σ one has that $(N_\sigma)_{i,j} = 1$ implies

$$(R_\sigma)_{i,k} = \begin{cases} (R_\sigma)_{i+1,k}, & \text{if } k < j; \\ (R_\sigma)_{i+1,k} + 1, & \text{if } k \geq j; \end{cases} \quad \text{and} \quad (R_\sigma)_{k,j} = \begin{cases} (R_\sigma)_{k,j-1}, & \text{if } k > i; \\ (R_\sigma)_{k,j-1} + 1, & \text{if } k \geq i. \end{cases} \tag{$*$}$$

This implies as well that $(N_\sigma)_{k,i} = (N_\sigma)_{j,k} = 0$ for any k so that

$$(R_\sigma)_{k,i} = (R_\sigma)_{k,i-1} \quad \text{and} \quad (R_\sigma)_{j,k} = (R_\sigma)_{j+1,k} \quad \text{for any } k\ : 1 \leq k \leq n. \tag{$**$}$$

Moreover, we get

Lemma. *Let* $\sigma \in \mathbf{S}_n^2$ *then* $(R_\sigma)_{i,j} = (R_\sigma)_{i+1,j} + 1 = (R_\sigma)_{i,j-1} + 1 = (R_\sigma)_{i+1,j-1} + 1$ *if and only if* $(i,j) \in \sigma$.

Proof.

If $(i,j) \in \sigma$ then $(N_\sigma)_{i,j} = 1$. This implies that $(R_\sigma)_{i,j} = (R_\sigma)_{i+1,j} + 1 = (R_\sigma)_{i,j-1} + 1$. Also $(N_\sigma)_{i,j} = 1$ implies $(N_\sigma)_{i,k} = (N_\sigma)_{t,j} = 0$ for all $k \neq j$ and $t \neq i$. Hence, in particular, $(R_\sigma)_{i+1,j} = (R_\sigma)_{i,j-1} = (R_\sigma)_{i+1,j-1}$.

On the other hand, assume that $(R_\sigma)_{i+1,j} = (R_\sigma)_{i,j-1} = (R_\sigma)_{i+1,j-1}$. This means by the form of N that $N_{i,k} = N_{t,j} = 0$ for any $k\ :\ 1 \leq k \leq j-1$ and for any $t\ :\ i+1 \leq t \leq n$. Then $(R_\sigma)_{i,j} = (R_\sigma)_{i+1,j} + 1$ forces $N_{i,j} = 1$.

■

Thus, by $(*), (**)$ and the lemma, any $R = R_\sigma$ for $\sigma \in \mathbf{S}_n^2$ has the following additional property

(iii) If $R_{i,j} = R_{i+1,j} + 1 = R_{i,j-1} + 1 = R_{i+1,j-1} + 1$ then
  (a) $R_{i,k} = R_{i+1,k}$ for any $k < j$ and $R_{i,k} = R_{i+1,k} + 1$ for any $k \geq j$;
  (b) $R_{k,j} = R_{k,j-1}$ for any $k > i$ and $R_{k,j} = R_{k,j-1} + 1$ for any $k \leq i$;
  (c) $R_{j,k} = R_{j+1,k}$ and $R_{k,i} = R_{k,i-1}$ for any $k\ :\ 1 \leq k \leq n$.

3.2. Let us study the structure of R_σ more explicitly.

Lemma. *Let* $R = R_\sigma$ *for some* $\sigma \in \mathbf{S}_n^2$. *For any* $i, j \ : \ 1 \le i < j \le n$ *only one of the following is possible*

(i) $R_{i,j} = R_{i+1,j-1}$ *and in that case* $R_{i+1,j} = R_{i,j-1} = R_{i,j}$;
(ii) $R_{i,j} = R_{i+1,j-1} + 2$ *and in that case* $R_{i+1,j} = R_{i,j-1} = R_{i+1,j-1} + 1$;
(iii) $R_{i,j} = R_{i+1,j-1} + 1$ *and* $R_{i+1,j} = R_{i,j-1} = R_{i+1,j-1}$;
(iv) $R_{i,j} = R_{i+1,j-1} + 1$ *and* $R_{i,j-1} = R_{i+1,j-1} + 1$, $R_{i+1,j} = R_{i,j}$;
(v) $R_{i,j} = R_{i+1,j-1} + 1$ *and* $R_{i+1,j} = R_{i+1,j-1} + 1$, $R_{i,j-1} = R_{i,j}$.

In each of the above cases R *can be presented graphically as follows:*

$$\text{(i)} \begin{array}{ccc} \cdots & p & p \\ \cdots & p & p \\ & \vdots & \vdots \end{array} \qquad \text{(ii)} \begin{array}{ccc} \cdots & p+1 & p+2 \\ \cdots & p & p+1 \\ & \vdots & \vdots \end{array}$$

$$\text{(iii)} \begin{array}{ccc} \cdots & p & p+1 \\ \cdots & p & p \\ & \vdots & \vdots \end{array} ; \quad \text{(iv)} \begin{array}{ccc} \cdots & p & p+1 \\ \cdots & p & p+1 \\ & \vdots & \vdots \end{array} , \quad \text{(v)} \begin{array}{ccc} \cdots & p+1 & p+1 \\ \cdots & p & p \\ & \vdots & \vdots \end{array}$$

Proof.

Consider some N_σ. The first case is simply equivalent to $N_{k,j} = N_{i,s} = 0$ for any $k \ge i$ and any $s \le j$. The second case is equivalent to the existence of $k > i$ and of $s < j$ such that $N_{k,j} = N_{i,s} = 1$. Since $R_{i,j}$ is the number of ones in the submatrix $N_{(i,j)}$ it is obvious that $R_{i,j} = R_{i+1,j-1} + 2$ in that case. The third case occurs if $N_{i,j} = 1$ and, thus, $N_{k,j} = N_{i,s} = 0$ for any $k > i$ and any $s < j$ so that $R_{i+1,j} = R_{i,j-1} = R_{i+1,j-1}$. The fourth case occurs if $N_{i,s} = 0$ for all $s \le j$ and $N_{k,j} = 1$ for some $k > i$. Finally, the fifth case occurs if $N_{k,j} = 0$ for all $k \ge i$ and $N_{i,s} = 1$ for some $s < j$.

■

3.3. Let $\mathbf{R}_n^2 \subset \mathbf{M}_n$ denote a subset of matrices satisfying properties (i)-(iii) of 3.1.

Proposition. $\mathbf{R}_n^2 = \{R_{N_\sigma}\}_{\sigma \in \mathbf{S}_n^2}$.

Proof.

We have seen already that $R_\sigma \in \mathbf{R}_n^2$ for all $\sigma \in \mathbf{S}_n^2$. On the other hand, consider some $R \in \mathbf{R}_n^2$ and let N be the matrix defined by

$$N_{i,j} = \begin{cases} 1, & \text{if } R_{i,j} = R_{i+1,j} + 1 = R_{i,j-1} + 1 = R_{i+1,j-1} + 1; \ (*) \\ 0, & \text{otherwise.} \end{cases}$$

By property (i) $N_{i,j} = 0$ for any $i, j \ : \ i \ge j$. Note that if $N_{i,j} = 1$ then $N_{i,k} = 0$ for any $k \ne j$. Indeed, $R_{i,k} = R_{i+1,k}$ for $k < j$ and $R_{i,k-1} = R_{i+1,k-1} + 1$ for $k > j$ by property iii(a) so that conditions $(*)$ are not satisfied for any (i, k) for which $k \ne j$. Exactly in the same way we get by iii(b) that $N_{k,j}=0$ for any $k\neq i$. Again, by iii(c) no $N_{j,k}$ and $N_{k,i}$ can satisfy conditions $(*)$, thus, $N_{j,k}=N_{k,i}=0$. Hence, $N=N_\sigma$ for σ defined by non-zero entries of N. It remains to show that $R_\sigma=R$. This is a straightforward corollary of (i) and (ii) of 3.1.

$\blacksquare$

For example, $\mathbf{R}_4^2$ contains 10 matrices

$$R_{(1,2)(3,4)}=\begin{pmatrix}0&1&1&2\\0&0&0&1\\0&0&0&1\\0&0&0&0\end{pmatrix};\ R_{(1,4)(2,3)}=\begin{pmatrix}0&0&1&2\\0&0&1&1\\0&0&0&0\\0&0&0&0\end{pmatrix};\ R_{(1,3)(2,4)}=\begin{pmatrix}0&0&1&2\\0&0&0&1\\0&0&0&0\\0&0&0&0\end{pmatrix};$$

$$R_{(1,2)}=\begin{pmatrix}0&1&1&1\\0&0&0&0\\0&0&0&0\\0&0&0&0\end{pmatrix};\quad R_{(1,3)}=\begin{pmatrix}0&0&1&1\\0&0&0&0\\0&0&0&0\\0&0&0&0\end{pmatrix};\quad R_{(1,4)}=\begin{pmatrix}0&0&0&1\\0&0&0&0\\0&0&0&0\\0&0&0&0\end{pmatrix};$$

$$R_{(2,3)}=\begin{pmatrix}0&0&1&1\\0&0&1&1\\0&0&0&0\\0&0&0&0\end{pmatrix};\quad R_{(2,4)}=\begin{pmatrix}0&0&0&1\\0&0&0&1\\0&0&0&0\\0&0&0&0\end{pmatrix};$$

$$R_{(3,4)}=\begin{pmatrix}0&0&0&1\\0&0&0&1\\0&0&0&1\\0&0&0&0\end{pmatrix};\quad R_{Id}=\begin{pmatrix}0&0&0&0\\0&0&0&0\\0&0&0&0\\0&0&0&0\end{pmatrix}.$$

3.4. Let $I_2\subset S(\mathfrak{n}^-)$ be the ideal generated by $I_2=\langle\sum_{k=j+1}^{i-1}x_{i,k}x_{k,j}\rangle_{j=1,i=j+1}^{j=n-1,i=n}$. Then $\mathcal{V}(I_2)=\mathcal{X}_2$.

Given $\sigma\in\mathbf{S}_n^2$ let $\mathcal{B}_\sigma$ be the corresponding $\mathbf{B}$ orbit in $\mathcal{X}_2$. $D_{\mathcal{B}_\sigma}$ (cf. 2.3) defines as in 2.9 the ideal $\mathcal{I}(\mathcal{B}_\sigma)$ or simply $\mathcal{I}(\sigma)$ of $S(\mathfrak{n}^-)$. which has two types of generating polynomials. The polynomials of the first type are quadratic (homogeneous) polynomials generating I_2 since $I_2\subset\mathcal{I}(\sigma)$. The generating polynomials of the second type in $\mathcal{I}_\sigma$ are determined by $\mathrm{Rank}\,(\pi_{i,j}(N_\sigma))$ that is by $(R_\sigma)_{i,j}$. We translate this fact into the set of minors of $\pi_{i,j}(N)$ of size $(R_\sigma)_{i,j}+1$. These polynomials are homogeneous of degree $(R_\sigma)_{i,j}+1$. Let us denote the ideal generated by the polynomials of the second type by $I_1(\sigma)$. Put $\mathcal{I}_{R_\sigma}:=\mathcal{I}_\sigma:=I_2+I_1(\sigma)$. Set $\mathcal{V}_{R_\sigma}:=\mathcal{V}_\sigma:=\mathcal{V}(\mathcal{I}_\sigma)$.

Note that $\pi_{i,j}(N_\sigma)$ provides new polynomials generating $I_1(\sigma)$ compared to those obtained from $\pi_{i+1,j}(N_\sigma)$ and $\pi_{i,j-1}(N_\sigma)$ only in the situation when $(R_\sigma)_{i,j}=(R_\sigma)_{i+1,j-1}$. Indeed, by lemma 2.9(ii) if $(R_\sigma)_{i,j}=(R_\sigma)_{i+1,j-1}+1$ then all the new generating polynomials obtained from $(R_\sigma)_{i,j}$ are in I_2.

For example, take $\sigma = (1,2)(3,4) \in \mathbf{S}_5^2$. One has

$$I_2 = \left\langle \begin{array}{c} x_{3,2}x_{2,1};\ x_{4,3}x_{3,2},\ x_{5,4}x_{4,3};\ x_{4,3}x_{3,1} + x_{4,2}x_{2,1}; \\ x_{5,4}x_{4,2} + x_{5,3}x_{3,2};\ x_{5,4}x_{4,1} + x_{5,3}x_{3,1} + x_{5,2}x_{2,1} \end{array} \right\rangle$$

and

$$R_{(1,2)(3,4)} = \begin{pmatrix} 0 & 1 & 1 & 2 & 2 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

So that

$$\begin{aligned} \mathcal{I}_{(1,2)(3,4)} &= \left\langle I_2; x_{3,2};\ x_{5,4};\ \begin{vmatrix} x_{4,2} & x_{4,3} \\ x_{5,2} & x_{5,3} \end{vmatrix} = x_{4,2}x_{5,3} - x_{4,3}x_{5,2} \right\rangle \\ &= \langle x_{3,2};\ x_{5,4};\ x_{4,3}x_{3,1} + x_{4,2}x_{2,1};\ x_{5,3}x_{3,1} + x_{5,2}x_{2,1};\ x_{4,2}x_{5,3} - x_{4,3}x_{5,2} \rangle \end{aligned}$$

Moreover, one can easily check that $\mathcal{I}_{(1,2)(3,4)}$ is a prime ideal.

By the definition of $\mathcal{I}_\sigma$ one can see at once that $N_\sigma \in \mathcal{V}_\sigma$. Hence,

Corollary. $\overline{\mathcal{B}}_\sigma \subset \mathcal{V}_\sigma$.

3.5. Let us define a partial order on $\mathbf{R}_n^2$. Given $R, R' \in \mathbf{R}_n^2$ we put $R \le R'$ if for any $1 \le i, j \le n$ one has $R_{i,j} \le R'_{i,j}$. Let us define the corresponding partial order on $\mathbf{S}_n^2$. For $\sigma, \sigma' \in \mathbf{S}_n^2$ we put $\sigma \le \sigma'$ if $R_\sigma \le R_{\sigma'}$.

The aim of this part is to prove

Theorem. *For any $\sigma \in \mathbf{S}_n^2$ one has*

$$\overline{\mathcal{B}}_\sigma = \mathcal{V}_\sigma = \coprod_{\sigma' \le \sigma} \mathcal{B}_{\sigma'}.$$

To do this we study the inclusion relation on $\{\mathcal{I}_\sigma\}_{\sigma \in \mathbf{S}_n^2}$ and show that it coincides with inverse inclusion relation on $\{\overline{\mathcal{B}}_\sigma\}_{\sigma \in \mathbf{S}_n^2}$. Then the result obtains as a corollary.

3.6. All the power rank conditions in the case of $\mathcal{X}_2$ are defined just by ranks of the minors of the lower-triangular matrix of indeterminants (and the common conditions coming from $X^2 = 0$). Hence $\sigma \le \sigma'$ (or $R_\sigma \le R_{\sigma'}$) $\iff \mathcal{I}_\sigma \supset \mathcal{I}_{\sigma'} \iff \mathcal{V}_\sigma \subset \mathcal{V}_{\sigma'}$.

Lemma. $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$ *implies* $\mathcal{V}_{\sigma'} \subset \mathcal{V}_\sigma$.

Proof.

The proof is by induction on n. It holds trivially for $n = 2$. Assume it holds for $n - 1$.

Assume $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$ for $\sigma', \sigma \in \mathbf{S}_n^2$.

(i) Since $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$ implies $\mathcal{O}_{\sigma'} \subset \overline{\mathcal{O}}_\sigma$, thus, by Gerstenhaber's result (cf. 2.8), $\operatorname{Rank} N_{\sigma'} \le \operatorname{Rank} N_\sigma$, that is, $(R_{\sigma'})_{1,n} \le (R_\sigma)_{1,n}$.

(ii) By 2.12 this implies as well that $\pi_{1,n-1}(\mathcal{B}_{\sigma'}) \subset \overline{\pi_{1,n-1}(\mathcal{B}_\sigma)}$. By the induction assumption this provides $R_{\pi_{1,n-1}(\sigma')} \leq R_{\pi_{1,n-1}(\sigma)}$.

(iii) Exactly in the same way we get $\pi_{2,n}(\mathcal{B}_{\sigma'}) \subset \overline{\pi_{2,n}(\mathcal{B}_\sigma)}$ and, thus, $R_{\pi_{2,n}(\sigma')} \leq R_{\pi_{2,n}(\sigma)}$.

By (ii) and (iii) we get $(R_{\sigma'})_{i,j} \leq (R_\sigma)_{i,j}$ for any $(i,j) \neq (1,n)$ and by (i) one has $(R_{\sigma'})_{1,n} \leq (R_\sigma)_{1,n}$. Thus, $R_{\sigma'} \leq R_\sigma$.

■

3.7. To show that $\mathcal{V}_{\sigma'} \subset \mathcal{V}_\sigma$ implies $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$ let us consider the following two subsets of $\{\sigma' \ : \sigma' < \sigma\}$:

(i) $D_1(\sigma)$ is the set of all σ' such that
  (a) $\sigma' < \sigma$;
  (b) $l(\sigma') = l(\sigma)$;
  (c) for any σ'' such that $\sigma \geq \sigma'' \geq \sigma'$ one has either $\sigma'' = \sigma$ or $\sigma'' = \sigma'$.

(ii) $D_2(\sigma)$ is the set of all σ' such that
  (a) $\sigma' < \sigma$;
  (b) $l(\sigma') < l(\sigma)$;
  (c) for any σ'' such that $l(\sigma'') < l(\sigma)$ and $\sigma > \sigma'' \geq \sigma'$ one has $\sigma'' = \sigma'$.

For any $\sigma' \in D_1(\sigma)$ one has $\mathcal{O}_{\sigma'} = \mathcal{O}_\sigma$. Moreover, $D_1(\sigma)$ is the set of the maximal elements of length $l(\sigma)$ which are smaller than σ with respect to the order defined in 3.5.

For every $\sigma' \in D_2(\sigma)$ one has $\overline{\mathcal{O}}_{\sigma'} \subsetneq \overline{\mathcal{O}}_\sigma$. Moreover $D_2(\sigma)$ is the set of the maximal elements of length less than $l(\sigma)$ which are smaller than σ with respect to the order defined in 3.5.

We begin with the construction of $D_2(\sigma)$. Then we construct $D_1(\sigma)$. Both constructions are straightforward but the proofs are rather long.

Recall from 2.5 that we write $\sigma = (i_1, j_1)(i_2, j_2)\ldots(i_k, j_k)$ where $i_s < j_s$ for any $s \ : 1 \leq s \leq k$ and $i_s < i_{s+1}$ for any $s \ : \ 1 \leq s < k$. Recall notation $(i,j) \in \sigma$, $I(\sigma)$ and $J(\sigma)$ from 2.13. Set

$$M(\sigma) := \{s \ : \ s = 1 \text{ or } s > 1 \text{ and } j_s > \max\{j_1, \ldots j_{s-1}\}\}.$$

The following lemma is straightforward.

Lemma. *If $s \in M(\sigma)$ then for any $(i,j) \in \sigma$ such that $(i,j) \neq (i_s, j_s)$ one has that either $i > i_s$ or $j < j_s$. In particular, fix $m := \max\{s \in M(\sigma)\}$ then $j_m = \max\{j \in J(\sigma)\}$.*

3.8. For any $s \ : \ 1 \leq s \leq k$ let $\sigma_s^- := (i_1, j_1)\ldots(i_{s-1}, j_{s-1})(i_{s+1}, j_{s+1})\ldots$ be the involution obtained by omitting (i_s, j_s). From the definition of $\leq$ one has $\sigma_s^- < \sigma$. Our aim is to show that $D_2(\sigma) = \{\sigma_s^- \mid s \in M(\sigma)\}$. We will first prove the

Lemma. $D_2(\sigma) \supseteq \{\sigma_s^- \mid s \in M(\sigma)\}$.

Proof.

Consider $\sigma = (i_1, j_1) \dots (i_k, j_k) \in \mathbf{S}_n^2$.

Assume $(i_1, j_1) = (1, n)$. Then $D_2(\sigma) = \{\sigma_1^- = (i_2, j_2) \dots (i_k, j_k)\}$ since for any $(i, j) \neq (1, n)$ one has $(R_\sigma)_{i,j} = (R_{\sigma_1^-})_{i,j}$ and $(R_{\sigma_1^-})_{1,n} = (R_\sigma)_{1,n} - 1$.

Now assume $(i_1, j_1) \neq (1, n)$. If $s \in M(\sigma)$ then

$$(R_{\sigma_s^-})_{p,q} = \begin{cases} (R_\sigma)_{p,q} - 1, & \text{if } p \leq i_s \text{ and } q \geq j_s; \\ (R_\sigma)_{p,q}, & \text{otherwise.} \end{cases}$$

Assume that there exists σ' such that $l(\sigma') < k$ and $R_\sigma > R_{\sigma'} \geq R_{\sigma_s^-}$. Since $(R_{\sigma_s^-})_{p,q} = (R_\sigma)_{p,q}$ whenever $p > i_s$ or $q < j_s$ we obtain $(R_{\sigma'})_{p,q} = (R_{\sigma_s^-})_{p,q} = (R_\sigma)_{p,q}$ whenever $p > i_s$ or $q < j_s$.

In particular, for any $(i, j) \in \sigma_s^-$ we have either $i > i_s$ or $j < j_s$ by lemma 3.7. Hence, $(R_{\sigma'})_{p,q} = (R_{\sigma_s^-})_{p,q} = (R_\sigma)_{p,q}$ for $(p, q) \in \{(i+1, j-1),\ (i, j-1),\ (i+1, j),\ (i, j)\}$. Further, since $(N_\sigma)_{i,j} = 1$ lemma 3.1 gives $(R_\sigma)_{i,j} = (R_\sigma)_{i+1,j-1} + 1 = (R_\sigma)_{i,j-1} + 1 = (R_\sigma)_{i+1,j} + 1$. Thus, also $(R_{\sigma'})_{i,j} = (R_{\sigma'})_{i+1,j-1} + 1 = (R_{\sigma'})_{i,j-1} + 1 = (R_{\sigma'})_{i+1,j} + 1$, which by lemma 3.1 gives $(i, j) \in \sigma'$ whenever $(i, j) \in \sigma_s^-$. Finally, since $l(\sigma') \leq k - 1$ we obtain $\sigma' = \sigma_s^-$.

■

3.9.

Proposition. $D_2(\sigma) = \{\sigma_s^- \mid s \in M(\sigma)\}$.

Proof.

By lemma 3.8 it remains to show that for any σ' such that $\sigma' < \sigma$ and $l(\sigma') < l(\sigma)$ there exists $s \in M(\sigma)$ such that $\sigma' \leq \sigma_s^-$.

The proof is by induction on n. It is trivially true for $\sigma \in \mathbf{S}_2^2$. Assume it is true for $\sigma \in \mathbf{S}_{n-1}^2$. Take $\sigma = (i_1, j_1) \dots (i_k, j_k) \in \mathbf{S}_n^2$ and $\sigma' = (i'_1, j'_1) \dots (i'_{k'}, j'_{k'}) \in \mathbf{S}_n^2$ such that $\sigma' < \sigma$ and $k' < k$. In the induction step we can consider recursion either to $\pi_{1,n-1}$ or to $\pi_{2,n}$. This involves the interplay between all possible $(R_{\sigma'})_{i,j}$ where $i = 1, 2,\ j = n-1, n$. The argument is broken into the five claims below. We formulate the assumptions of each claim both in terms of $J(\sigma'),\ J(\sigma)$ (resp. $I(\sigma'),\ I(\sigma)$) and in terms of $(R_{\sigma'})_{1,n-1},\ (R_\sigma)_{1,n-1}$ (resp. $(R_{\sigma'})_{2,n},\ (R_\sigma)_{2,n}$) for greater clarity. The conclusion of each claim is the existence of $s \in M(\sigma)$ such that $\sigma' \leq \sigma_s^-$.

Claim 1. *Assume that $n \notin J(\sigma'), J(\sigma)$ (resp. $1 \notin I(\sigma'), I(\sigma)$). This is equivalent to $(R_{\sigma'})_{1,n-1} = (R_{\sigma'})_{1,n}$ and $(R_\sigma)_{1,n-1} = (R_\sigma)_{1,n}$ (resp. $(R_{\sigma'})_{2,n} = (R_{\sigma'})_{1,n}$ and $(R_\sigma)_{2,n} = (R_\sigma)_{1,n}$.) Then there exists $s \in M(\sigma)$ such that $\sigma' \leq \sigma_s^-$.*

Proof.

By symmetry about the anti-diagonal it is enough to consider the case $n \notin J(\sigma'), J(\sigma)$ that is $(R_{\sigma'})_{1,n-1} = (R_{\sigma'})_{1,n} = k'$ and $(R_\sigma)_{1,n-1} = (R_\sigma)_{1,n} = k$. In that case $\pi_{1,n-1}(\sigma') = \sigma'$ and $\pi_{1,n-1}(\sigma) = \sigma$. By the induction hypothesis there exists $s \in M(\sigma)$ such that $\sigma > \sigma_s^- \geq \sigma'$. Considering σ, σ_s^- and σ' as elements of $\mathbf{S}_n^2$ we get the corresponding rank matrices by adding an $n-$th row with zero entries and an $n-$th column with entries equal to those in the $(n-1)$-th column. Thus, $\sigma' \leq \sigma_s^-$ as elements of $\mathbf{S}_n^2$.

■

Claim 2. *Assume that* $n \notin J(\sigma')$ *and* $n \in J(\sigma)$ *(resp.* $1 \notin I(\sigma')$ *and* $1 \in I(\sigma)$*). This is equivalent to* $(R_{\sigma'})_{1,n-1} = (R_{\sigma'})_{1,n}$ *(resp.* $(R_{\sigma'})_{2,n} = (R_{\sigma'})_{1,n}$*) and* $(R_\sigma)_{1,n-1} = (R_\sigma)_{1,n} - 1$ *(resp.* $(R_\sigma)_{2,n} = (R_\sigma)_{1,n} - 1$*). Then* $\sigma' \leq \sigma_m^-$ *where* $m = \max\{s \in M(\sigma)\}$ *(resp.*$\sigma' \leq \sigma_1^-$*).*

Proof.

Again, by symmetry about the anti-diagonal it is enough to consider the case $(R_{\sigma'})_{1,n-1} = (R_{\sigma'})_{1,n}$ and $(R_\sigma)_{1,n-1} = (R_\sigma)_{1,n} - 1$. In that case $j_m = n$. Let us show that $\sigma' \leq \sigma_m^-$. Indeed, $(R_{\sigma_m^-})_{i,j} = (R_\sigma)_{i,j}$ whenever $j < n$ so that $(R_{\sigma'})_{i,j} \leq (R_{\sigma_m^-})_{i,j}$ for such pairs. Since $n \notin J(\sigma')$ one has $(R_{\sigma'})_{i,n} = (R_{\sigma'})_{i,n-1} \leq (R_{\sigma_m^-})_{i,n-1} = (R_{\sigma_m^-})_{i,n}$ for any i. Thus $\sigma' \leq \sigma_m^-$.

■

We are left with the case $1 \in I(\sigma')$ and $n \in J(\sigma')$. We have two subcases which we consider in three claims below.

Claim 3. *Assume that* $(1, n) \in \sigma'$*. This is equivalent to* $(R_{\sigma'})_{2,n-1} = (R_{\sigma'})_{1,n-1} = (R_{\sigma'})_{2,n} = k' - 1$*. Then there exists* $s \in M(\sigma)$ *such that* $\sigma' < \sigma_s^-$*.*

Proof.

If $1 \in I(\sigma)$ then $\sigma' \leq \sigma_1^-$ since

(i) $(R_{\sigma_1^-})_{i,j} = (R_\sigma)_{i,j} \geq (R_{\sigma'})_{i,j}$ whenever $i > 1$;
(ii) $(R_{\sigma'})_{1,j} = (R_{\sigma'})_{2,j} \leq (R_\sigma)_{2,j} = (R_{\sigma_1^-})_{1,j}$ whenever $j < n$;
(iii) $(R_{\sigma'})_{1,n} = k' \leq k - 1 = (R_{\sigma_1^-})_{1,n}$;

If $1 \notin I(\sigma)$ then $\pi_{2,n}(\sigma) = \sigma$ and $\pi_{2,n}(\sigma') = (\sigma')_1^-$. By the induction hypothesis there exists $s \in M(\sigma)$ such that $\sigma_s^- \geq (\sigma')_1^-$. Let us show that $\sigma_s^- \geq \sigma'$. Indeed,

(i) $(R_{\sigma_s^-})_{i,j} \geq (R_{(\sigma')_1^-}) = (R_{\sigma'})_{i,j}$ whenever $i > 1$;
(ii) $(R_{\sigma'})_{1,j} = (R_{\sigma'})_{2,j} \leq (R_{\sigma_s^-})_{2,j} = (R_{\sigma_s^-})_{1,j}$ whenever $j < n$;
(iii) $(R_{\sigma'})_{1,n} = k' \leq k - 1 = (R_{\sigma_s^-})_{1,n}$;

Note that in both cases the inequality is strict since in both cases $(R_{\sigma_s^-})_{2,n} = k - 1 > k' - 1 = (R_{\sigma'})_{2,n}$.

■

The remaining case is when $1 \in I(\sigma')$ and $n \in J(\sigma')$, but $(1,n) \notin \sigma'$. We consider two cases according to whether n is or is not contained in $J(\sigma)$.

Claim 4. *Assume that* $1 \in I(\sigma')$ *and* $n \in J(\sigma')$ *but* $(1,n) \notin \sigma'$. *This is equivalent to* $(R_{\sigma'})_{2,n-1} = k'-2$, $(R_{\sigma'})_{1,n-1} = (R_{\sigma'})_{2,n} = k'-1$. *Assume also that* $n \notin J(\sigma)$. *This is equivalent to* $(R_\sigma)_{1,n-1} = (R_\sigma)_{1,n}$. *Then there exists* $s \in M(\sigma)$ *such that* $\sigma' < \sigma_s^-$.

Proof.

If $n \notin J(\sigma)$ then $\pi_{1,n-1}(\sigma) = \sigma$ and $l(\pi_{1,n-1}(\sigma)) - l(\pi_{1,n-1}(\sigma')) = k-(k'-1) \geq 2$. This inequality permits us to use the induction hypothesis twice. Thus, there exists $s \in M(\sigma)$ and $t \in M(\sigma_s^-)$ such that $\pi_{1,n-1}(\sigma') \leq (\sigma_s^-)_t^- < \sigma_s^-$. Let i_m be $\min\{i_s, i_t\}$ then $m \in M(\sigma)$ and $(R_{(\sigma_s^-)_t^-})_{i,j} \leq (R_{\sigma_m^-})_{i,j}$ for all possible (i,j). In particular, $(R_{\sigma'})_{i,j} \leq (R_{\sigma_m^-})_{i,j}$ whenever $j < n$. Considering $\sigma_m^-, (\sigma_s^-)_t^-$ as elements of $\mathbf{S}_n^2$ we get

$$(R_{\sigma_m^-})_{l,n} = \begin{cases} (R_\sigma)_{l,n}, & \text{if } l > i_m; \\ (R_{(\sigma_s^-)_t^-})_{l,n-1} + 1, & \text{if } l \leq i_m. \end{cases}$$

On the other hand $(R_{\sigma'})_{l,n} \leq (R_{\sigma'})_{l,n-1} + 1 \leq (R_{(\sigma_s^-)_t^-})_{l,n-1} + 1$ for any l by definition of $(\sigma_s^-)_t^-$ and the construction of the rank matrices. Thus, for $l \leq i_m$ we get $(R_{\sigma'})_{l,n} \leq (R_{\sigma_m^-})_{l,n} = (R_{(\sigma_s^-)_t^-})_{l,n-1} + 1$. Finally, for $l > i_m$ we get $(R_{\sigma'})_{l,n} \leq (R_{\sigma_m^-})_{l,n} = (R_\sigma)_{l,n}$. Altogether, this gives $\sigma' < \sigma_m^-$. Note that the inequality is strict since $(R_{\sigma_m^-})_{1,n-1} = k-1 > (R_{\sigma'})_{1,n-1} = k'-1$.

■

Claim 5. *Assume that* $1 \in I(\sigma')$ *and* $n \in J(\sigma')$ *but* $(1,n) \notin \sigma'$. *This is equivalent to* $(R_{\sigma'})_{2,n-1} = k'-2$, $(R_{\sigma'})_{1,n-1} = (R_{\sigma'})_{2,n} = k'-1$. *Assume also that* $n \in J(\sigma)$. *This is equivalent to* $(R_\sigma)_{1,n-1} = (R_\sigma)_{1,n} - 1$. *Then there exists* $s \in M(\sigma)$ *such that* $\sigma' \leq \sigma_s^-$.

Proof.

Again, consider $\pi_{1,n-1}(\sigma), \pi_{1,n-1}(\sigma')$. Note that $\pi_{1,n-1}(\sigma) > \pi_{1,n-1}(\sigma')$. We also have $l(\pi_{1,n-1}(\sigma)) = k-1 > l(\pi_{1,n-1}(\sigma')) = k'-1$. Thus, by induction hypothesis there exists $s \in M(\pi_{1,n-1}(\sigma))$ such that $\pi_{1,n-1}(\sigma') \leq (\pi_{1,n-1}(\sigma))_s^-$. Let $r = \max\{t \in M(\sigma)\}$ then $j_r = n$. Let $m = \min\{i_s, i_r\}$. Again, $m \in M(\sigma)$ by lemma 3.7 and $(R_{\sigma'})_{i,j} \leq (R_{\sigma_m^-})$ whenever $j < n$. Note also

$$(R_{\sigma_m^-})_{l,n} = \begin{cases} (R_\sigma)_{l,n}, & \text{if } l > i_m; \\ (R_{\sigma_s^-})_{l,n-1} + 1, & \text{if } l \leq i_m. \end{cases}$$

Thus, for $l > i_m$ one has $(R_{\sigma'})_{l,n} \leq (R_\sigma)_{l,n} = (R_{\sigma_m^-})_{l,n}$ and for $l \leq i_m$ one has $(R_\sigma)_{l,n} \leq (R_\sigma)_{l,n-1} + 1 \leq (R_{\sigma_s^-})_{l,n-1} + 1 = (R_{\sigma_m^-})_{l,n}$. Altogether this gives $\sigma' \leq \sigma_m^-$.

■

In claims 1-5 we have considered all the possible cases described in lemma 3.2. Therefore, the proof is completed.

■

For example, take $\sigma=(1,6)(2,3)(4,5)(7,8)(9,12)(10,11)$. Then

$$D_2(\sigma)=\left\{\begin{array}{c}(2,3)(4,5)(7,8)(9,12)(10,11),\ (1,6)(2,3)(4,5)(9,12)(10,11),\\ (1,6)(2,3)(4,5)(7,8)(10,11)\end{array}\right\}$$

3.10. Now we construct $D_1(\sigma)$. It consists of four types of elements. In the next four subsections we define these types and show that an element of the given type is in $D_1(\sigma)$. Then in subsection 3.15 we show that $D_1(\sigma)$ consists of these four types of elements only.

In the next five sections we set $\sigma=(i_1,j_1)\ldots(i_k,j_k)$ where $i_s<i_{s+1}$ for any $s:\ 1\leq s<k$ and $i_s<j_s$ for any $s:\ 1\leq s\leq k$.

Lemma. . *Let $\sigma,\sigma',\sigma''\in \mathbf{S}_n^2$ be such that $\sigma\geq\sigma'\geq\sigma''$ and $l(\sigma)=l(\sigma')=l(\sigma'')$.*

- (i) *If $I(\sigma)=I(\sigma'')$ then $I(\sigma')=I(\sigma)$.*
- (ii) *If $J(\sigma)=J(\sigma'')$ then $J(\sigma')=J(\sigma)$.*

Proof.

(i) Assume that $I(\sigma)=I(\sigma'')$. Then by definition of a rank matrix $(R_\sigma)_{i,n}=(R_{\sigma''})_{i,n}$ for all i. Since $(R_\sigma)_{i,n}\leq(R_{\sigma'})_{i,n}\leq(R_{\sigma''})_{i,n}$ this forces $(R_\sigma)_{i,n}=(R_{\sigma'})_{i,n}$ for all i. Thus $(R_{\sigma'})_{i,n}=(R_{\sigma'})_{i+1,n}+1\iff(R_\sigma)_{i,n}=(R_\sigma)_{i+1,n}+1$. Hence, $i\in I(\sigma)$ iff $i\in I(\sigma')$.

(ii) is obtained exactly in the same way.

■

3.11. Let $(i_s,j_s)\in\sigma$. Suppose there exists $m<i_s$ satisfying both

- (i) $m\notin I(\sigma)\cup J(\sigma)$;
- (ii) Either $m=i_s-1$ or for any $t\ :\ m<t<i_s$ one has $t\in I(\pi_{1,j_s}(\sigma))\cup J(\pi_{1,j_s}(\sigma))$.

Then define $\sigma_{i_s\uparrow}$ to be obtained from σ by just changing (i_s,j_s) to (m,j_s). By the definition such an m is unique, if it exists. If such an m does not exist put $\sigma_{i_s\uparrow}:=\emptyset$.

For example, take $\sigma=(2,5)(3,4)(7,9)(8,10)$. Then $\sigma_{2\uparrow}=(1,5)(3,4)(7,9)(8,10)$, $\sigma_{3\uparrow}=\emptyset$, $\sigma_{7\uparrow}=(2,5)(3,4)(6,9)(8,10)$ and $\sigma_{8\uparrow}=(2,5)(3,4)(7,9)(6,10)$.

If $\sigma_{i_s\uparrow}\neq\emptyset$ then

$$(R_{\sigma_{i_s}})_{i,j}=\begin{cases}(R_\sigma)_{i,j}-1, & \text{if } m<i\leq i_s \text{ and } j\geq j_s;\\ (R_\sigma)_{i,j}, & \text{otherwise;}\end{cases}\qquad(*)$$

Thus, $\sigma_{i_s\uparrow}<\sigma$.

Lemma. *Take $\sigma=(i_1,j_1)\ldots(i_k,j_k)$. Let $s\ :\ 1\leq s\leq k$ be such that $\sigma_{i_s\uparrow}\neq\emptyset$. Then $\sigma_{i_s\uparrow}\in D_1(\sigma)$.*

Proof.

The proof is by induction on n. The minimal n which permits such a situation to occur is $n = 3$. In $\mathbf{S}_3^2$ there is a unique case of this type: $\sigma = (2,3)$ and $\sigma_{2\uparrow} = (1,3)$. It is obvious that $\sigma_{2\uparrow} \in D_1(\sigma)$. Assume the claim is true for $n-1$.

Suppose $\sigma_{i_s\uparrow} \neq \emptyset$ for some s. Let $\sigma' = (i'_1, j'_1)\ldots(i'_k, j'_k)$ be such that $\sigma \geq \sigma' \geq \sigma_{i_s\uparrow}$. We must show that either $\sigma' = \sigma$ or $\sigma' = \sigma_{i_s\uparrow}$.

Note that by definition $J(\sigma) = J(\sigma_{i_s\uparrow})$. By lemma 3.10 this implies $J(\sigma') = J(\sigma)$. Set $J := J(\sigma)$.

Assume first that $n \notin J$. Passing to $\pi_{1,n-1}$ we get $\pi_{1,n-1}(\sigma) = \sigma$, $\pi_{1,n-1}(\sigma') = \sigma'$ and $\pi_{1,n-1}(\sigma_{i_s\uparrow}) = \sigma_{i_s\uparrow}$ so that by the induction hypothesis either $\sigma' = \sigma$ or $\sigma' = \sigma_{i_s\uparrow}$.

Now assume that $n \in J$. We have to consider two cases: either $j_s = n$ or $j_s < n$.

- (i) Assume first that $j_s = n$. Then $\pi_{1,n-1}(\sigma_{i_s\uparrow}) = \pi_{1,n-1}(\sigma)$ so that $(R_\sigma)_{i,j} = (R_{\sigma_{i_s\uparrow}})_{i,j} = (R_{\sigma'})_{i,j}$ whenever $j < n$. This is equivalent to $(i_r, j_r) \in \sigma$ if and only if $(i_r, j_r) \in \sigma_{i_s\uparrow}$ and thus also if and only if $(i_r, j_r) \in \sigma'$ whenever $r \neq s$. Suppose $(i'_s, n) \in \sigma'$. Note that $I(\sigma) \setminus \{i_s\} = I(\sigma_{i_s\uparrow}) \setminus \{m\} = I(\sigma') \setminus \{i'_s\}$. Denote this common set by $\hat{I} := I(\sigma) \setminus \{i_s\}$.

  By $(*)$ and the structure of the rank matrices $(R_{\sigma_{i_s\uparrow}})_{l,n} = (R_\sigma)_{l,n} = (R_\sigma)_{l,n-1}$ whenever $l > i_s$ and $(R_{\sigma_{i_s\uparrow}})_{l,n} = (R_\sigma)_{l,n} = (R_\sigma)_{l,n-1} + 1$ whenever $l \leq m$. These equalities force $(R_{\sigma'})_{l,n} = (R_{\sigma'})_{l,n-1}$ whenever $l > i_s$ and $(R_{\sigma'})_{l,n} = (R_{\sigma'})_{l,n-1} + 1$ whenever $l \leq m$. Thus, by the structure of the rank matrices we get $m \leq i'_s \leq i_s$. Further, by the definition of $\sigma_{i_s\uparrow}$ for any $t\ :\ m < t < i_s$ one has $t \in \hat{I} \cup J$ so that either $i'_s = i_s$ (that is $\sigma' = \sigma$) or $i'_s = m$ (that is $\sigma' = \sigma_{i_s\uparrow}$).
- (ii) Now assume that $j_s < n$.
  - (a) Suppose $(i_r, n) \in \sigma$ (then $(i_r, n) \in \sigma_{i_s\uparrow}$ as well) and $(i'_r, n) \in \sigma'$. Let us show that $i'_r = i_r$. Indeed, by definition of $\sigma_{i_s\uparrow}$ one has either $i_r > i_s$ or $i_r < m$. In both cases one has by $(*)$ that $(R_{\sigma_{i_s\uparrow}})_{i_r,n-1} = (R_\sigma)_{i_r,n-1}$ and $(R_{\sigma_{i_s\uparrow}})_{i_r,n} = (R_\sigma)_{i_r,n} = (R_\sigma)_{i_r,n-1}+1$. One also has $(R_{\sigma_{i_s\uparrow}})_{i_r+1,n} = (R_{\sigma_{i_s\uparrow}})_{i_r,n} - 1 = (R_\sigma)_{i_r,n} - 1 = (R_\sigma)_{i_r+1,n}$. These equalities force $(R_{\sigma'})_{i_r+1,n} = (R_{\sigma'})_{i_r,n-1} = (R_{\sigma'})_{i_r,n} - 1$. By the structure of the rank matrices this implies $i'_r = i_r$.
  - (b) Since $j_s < n$ one has $\pi_{1,n-1}(\sigma_{i_s\uparrow}) = (\pi_{1,n-1}(\sigma))_{i_s\uparrow}$. Thus, by the induction hypothesis either $\pi_{1,n-1}(\sigma') = \pi_{1,n-1}(\sigma)$ or $\pi_{1,n-1}(\sigma') = \pi_{1,n-1}(\sigma_{i_s\uparrow})$. By (a) this implies that either $\sigma' = \sigma$ or $\sigma' = \sigma_{i_s\uparrow}$.

■

3.12. Again, let $(i_s, j_s) \in \sigma$. Suppose there exists $m > j_s$ satisfying both

- (i) $m \notin I(\sigma) \cup J(\sigma)$;
- (ii) Either $m = j_s + 1$ or for any $t\ :\ j_s < t < m$ one has $t \in I(\pi_{i_s,n}(\sigma)) \cup J(\pi_{i_s,n}(\sigma))$.

Then define $\sigma_{j_s\to}$ to be obtained from σ just by changing (i_s, j_s) to (i_s, m). Again, such an m is unique, if it exists. If such an m does not exist put $\sigma_{j_s\to} := \emptyset$.

For example, take $\sigma = (2,5)(3,4)(7,9)(8,10) \in \mathbf{S}^2_{11}$. Then $\sigma_{5\to} = (2,6)(3,4)(7,9)(8,10)$, $\sigma_{4\to} = \emptyset$, $\sigma_{9\to} = (2,5)(3,4)(7,11)(8,10)$ and $\sigma_{10\to} = (2,5)(3,4)(7,9)(8,11)$.

By symmetry of the rank matrix about the anti-diagonal if $\sigma_{j_s\to} \neq \emptyset$ then $\sigma_{j_s\to} < \sigma$ and

$$(R_{\sigma_{j_s\to}})_{i,j} = \begin{cases} (R_\sigma)_{i,j} - 1, & \text{if } j_s \le j < m \text{ and } i \le i_s; \\ (R_\sigma)_{i,j}, & \text{otherwise;} \end{cases} \tag{$*'$}$$

Exactly as in 3.11 we get the

Lemma. *Take $\sigma = (i_1, j_1) \dots (i_k, j_k)$. Let $s \ : \ 1 \le s \le k$ be such that $\sigma_{j_s\to} \neq \emptyset$. Then $\sigma_{j_s\to} \in D_1$.*

3.13. Consider $(i_s, j_s) \in \sigma$ for $s \ge 2$ and suppose there exists $j_r < i_s$ satisfying one of the conditions: either $j_r = i_s - 1$ or for any $t \ : \ j_r < t < i_s$ one has $t \in I(\pi_{i_r, j_s}(\sigma)) \cup J(\pi_{i_r, j_s}(\sigma))$. Then define σ_{j_r, i_s} to be obtained from σ just by changing the cycles $(i_r, j_r),\ (i_s, j_s)$ to the cycles $(i_r, i_s),\ (j_r, j_s)$.

Note that for a given i_s there can exist a few possible j_r. Let us denote the set of all possible σ_{j_r, i_s} for a given i_s by $C_{i_s\uparrow\to}(\sigma)$. If there is no $r < s$ such that j_r satisfies conditions put $C_{i_s\uparrow\to}(\sigma) := \emptyset$.

For example, take $\sigma = (1,3)(2,4)(5,8)(6,7)$. Then $C_{2\uparrow\to}(\sigma) = \emptyset$, $C_{5\uparrow\to}(\sigma) = \{\sigma_{4,5} = (1,3)(2,5)(4,8)(6,7), \sigma_{3,5} = (1,5)(2,4)(3,8)(6,7)\}$ and $C_{6\uparrow\to} = \emptyset$.

When σ_{j_r, i_s} is defined one has

$$(R_{\sigma_{j_r, i_s}})_{i,j} = \begin{cases} (R_\sigma)_{i,j} - 1, & \text{if } i \le i_r \text{ and } j_r \le j < i_s; \\ (R_\sigma)_{i,j} - 1, & \text{if } i_s \le i < j_r \text{ and } j \ge j_s; \\ (R_\sigma)_{i,j}, & \text{otherwise;} \end{cases} \tag{$**$}$$

Thus, $\sigma_{j_r, i_s} < \sigma$.

Lemma. *Take $\sigma = (i_1, j_1) \dots (i_k, j_k)$. Let $s \ : \ 2 \le s \le k$ and $r \ : \ r < s$ be such that σ_{j_r, i_s} exists. Then $\sigma_{j_r, i_s} \in D_1$.*

Proof.

The proof is by induction on n. The minimal n which permits such a situation to occur is $n = 4$. In $\mathbf{S}^2_4$ there is a unique case of this type: $\sigma = (1,2)(3,4)$ and $\sigma_{2,3} = (1,3)(2,4)$. It is obvious that $\sigma_{2,3} \in D_1(\sigma)$. Assume the claim is true for $n - 1$.

Suppose $\sigma_{j_r, i_s} \in C_{i_s\uparrow\to}(\sigma)$. Let $\sigma' = (i'_1, j'_1) \dots (i'_k, j'_k)$ be such that $\sigma_{j_r, i_s} \le \sigma' \le \sigma$. We must show either $\sigma' = \sigma$ or $\sigma' = \sigma_{j_r, i_s}$.

Note that $I(\sigma)\cup J(\sigma) = I(\sigma_{j_r,i_s})\cup J(\sigma_{j_r,i_s})$. In particular, one has $1 \in I(\sigma)$ if and only if $1 \in I(\sigma_{j_r,i_s})$. By $(**)$ $(R_\sigma)_{2,n} = (R_{\sigma_{j_r,i_s}})_{2,n}$, thus, $(R_{\sigma'})_{2,n} = (R_\sigma)_{2,n}$. Also, $(R_\sigma)_{2,n} = k-1$ if and only if $1 \in I(\sigma)$. Hence, $1 \in I(\sigma') \iff 1 \in I(\sigma)$.

Assume first that $1 \notin I(\sigma)$. Passing to $\pi_{2,n}$ we get $\pi_{2,n}(\sigma) = \sigma$, $\pi_{2,n}(\sigma') = \sigma'$ and $\pi_{2,n}(\sigma_{j_r,i_s}) = \sigma_{j_r,i_s}$ so that by the induction hypothesis either $\sigma' = \sigma$ or $\sigma' = \sigma_{j_r,i_s}$.

Now, let $1 \in I(\sigma)$.

(i) Assume first that $i_r > 1$. Then $\pi_{2,n}(\sigma_{j_r,i_s}) = (\pi_{2,n}(\sigma))_{j_r,i_s}$, thus, by the induction hypothesis $\pi_{2,n}(\sigma_{j_r,i_s}) \in D_1((\pi_{2,n}(\sigma)))$ so that either $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma)$ or $\pi_{2,n}(\sigma') = (\pi_{2,n}(\sigma))_{j_r,i_s}$.

  (a) If $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma)$ then $\sigma' < \sigma$ only if $j'_1 > j_1$. In that case $(R_{\sigma'})_{1,j_1} = (R_\sigma)_{1,j_1} - 1$. On the other hand, since $j_1 \notin [j_r, i_s]$ one has by $(**)$ that $(R_{\sigma_{j_r,i_s}})_{1,j_1} = (R_\sigma)_{1,j_1}$, which contradicts the condition $\sigma' \geq \sigma_{j_r,i_s}$. Thus, $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma)$ implies $\sigma' = \sigma$.

  (b) If $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma_{j_r,i_s})$ then $\sigma' > \sigma_{j_r,i_s}$ only if $j'_1 < j_1$. In that case $(R_{\sigma'})_{1,j'_1} = (R_{\sigma_{j_r,i_s}})_{1,j'_1} + 1$. On the other hand since $\pi_{2,n}(\sigma') = (\pi_{2,n}(\sigma))_{j_r,i_s}$ one has $j'_1 \notin [j_r, i_s]$ so that by $(**)$ $(R_{\sigma_{j_r,i_s}})_{1,j'_1} = (R_\sigma)_{1,j'_1}$ which contradicts the condition $\sigma' \leq \sigma$. Thus, $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma_{j_r,i_s})$ implies $\sigma' = \sigma_{j_r,i_s}$.

(ii) Assume that $i_r = 1$. In that case $r = 1$ and $\pi_{2,n}(\sigma_{j_1,i_s}) = (\pi_{2,n}(\sigma))_{i_s\uparrow}$. Thus, by Lemma 3.11 either $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma)$ or $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma_{j_1,i_s})$.

Assume first that $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma)$. Then $\sigma' < \sigma$ only if $j'_1 > j_1$. By the conditions on the construction of σ_{j_1,i_s} this implies $j'_1 > i_s$ so that by $(**)$ $(R_{\sigma_{j_1,i_s}})_{1,i_s} = (R_{\sigma'})_{1,i_s} + 1$ which contradicts $\sigma' \geq \sigma_{j_1,i_s}$. Thus, $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma)$ implies $\sigma' = \sigma$.

Now assume that $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma_{j_1,i_s})$. Then $\sigma' > \sigma_{j_1,i_s}$ only if $j'_1 < i_s$. Again, by the conditions on construction of σ_{j_1,i_s} this implies $j'_1 < j_1$ so that by $(**)$ $(R_{\sigma'})_{1,j'_1} = (R_\sigma)_{1,j'_1} + 1$ which contradicts $\sigma' \leq \sigma$. Thus, $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma_{j_1,i_s})$ implies $\sigma' = \sigma_{j_1,i_s}$.

■

3.14. In the construction of the last type we use the notion $M(\sigma)$ from 3.7. For $(i_s, j_s) \in \sigma$ consider $\pi_{i_s+1,j_s-1}(\sigma)$ and let $M_{[i_s,j_s]}(\sigma) := M(\pi_{i_s+1,j_s-1}(\sigma))$. If $M_{[i_s,j_s]}(\sigma) \neq \emptyset$ then for $t \in M_{[i_s,j_s]}(\sigma)$ define σ_{i_s,i_t} to be obtained from σ just by changing the cycles (i_s, j_s), (i_t, j_t) to the cycles (i_s, j_t), (i_t, j_s). If $M_{[i_s,j_s]}(\sigma) = \emptyset$ put $C_{i_s\uparrow\downarrow}(\sigma) := \emptyset$ otherwise put $C_{i_s\uparrow\downarrow}(\sigma) := \{\sigma_{i_s,i_t} \ : \ t \in M_{[i_s,j_s]}(\sigma)\}$.

For example, take $\sigma = (1,8)(2,5)(3,4)(6,7)$. Then

$$\begin{array}{rcl} C_{1\uparrow\downarrow}(\sigma) & = & \{\sigma_{1,2} = (1,5)(2,8)(3,4)(6,7),\ \sigma_{1,6} = (1,7)(2,5)(3,4)(6,8)\}, \\ C_{2\uparrow\downarrow}(\sigma) & = & \{\sigma_{2,3} = (1,8)(2,4)(3,5)(6,7)\}, \\ C_{3\uparrow\downarrow}(\sigma) & = & C_{6\uparrow\downarrow}(\sigma) = \emptyset. \end{array}$$

When σ_{i_s,i_t} is defined one has,

$$(R_{\sigma_{i_s,i_t}})_{i,j} = \begin{cases} (R_\sigma)_{i,j} - 1, & \text{if } i_t \leq i < i_s \text{ and } j_t \leq j < j_s; \\ (R_\sigma)_{i,j}, & \text{otherwise.} \end{cases} \qquad (***)$$

Thus, $\sigma_{i_s,i_t} < \sigma$.

In what follows we also need the following form of $(***)$:

$$(R_{\sigma_{i_s,i_t}})_{i,j} = \begin{cases} (R_{\sigma_t^-})_{i,j}, & \text{if } i > i_s \text{ and } j < j_s; \\ (R_\sigma)_{i,j}, & \text{otherwise.} \end{cases} \qquad (***')$$

Lemma. *Take $\sigma = (i_1, j_1) \dots (i_k, j_k)$. Let $s \ : \ 1 \leq s < k$ and $t \ : \ t > s$ be such that σ_{i_s,i_t} exists. Then $\sigma_{i_s,i_t} \in D_1$.*

Proof.

The proof is by induction on n. The minimal n which permits such situation to occur is $n = 4$. In $\mathbf{S}_4^2$ there is a unique case of this type: $\sigma = (1,4)(2,3)$ and $\sigma_{1,2} = (1,3)(2,4)$. It is obvious that $\sigma_{1,2} \in D_1(\sigma)$. Assume the claim is true for $n - 1$.

Suppose $\sigma_{i_s,i_t} \in C_{i_s \uparrow\downarrow}(\sigma)$ for some s. Let $\sigma' = (i'_1, j'_1) \dots (i'_k, j'_k)$ be such that $\sigma_{i_s,i_t} \leq \sigma' \leq \sigma$. We must show that either $\sigma' = \sigma$ or $\sigma' = \sigma_{i_s,i_t}$.

Note that $I(\sigma) = I(\sigma_{i_s,i_t})$ and $J(\sigma) = J(\sigma_{i_s,i_t})$. Thus, by lemma 3.10 $I(\sigma') = I(\sigma)$ and $J(\sigma') = J(\sigma)$. We will denote $I := I(\sigma)$ and $J := J(\sigma)$.

Assume first that $1 \notin I(\sigma)$. Passing to $\pi_{2,n}$ we get $\pi_{2,n}(\sigma) = \sigma$, $\pi_{2,n}(\sigma') = \sigma'$ and $\pi_{2,n}(\sigma_{i_s,i_t}) = \sigma_{i_s,i_t}$ so that by the induction hypothesis either $\sigma' = \sigma$ or $\sigma' = \sigma_{i_s,i_t}$.

Now, let $1 \in I(\sigma)$.

(i) Assume first that $s > 1$. Then $\pi_{2,n}(\sigma_{i_s,i_t}) = (\pi_{2,n}(\sigma))_{i_s,i_t}$, thus, by induction hypothesis $\pi_{2,n}(\sigma_{i_s,i_t}) \in D_1((\pi_{2,n}(\sigma)))$ so that either $\pi_{2,n}(\sigma') = \pi_{2,n}(\sigma)$ or $\pi_{2,n}(\sigma') = (\pi_{2,n}(\sigma))_{i_s,i_t}$. In both cases $i'_1 = 1$ and $j'_1 = j_1$ since $I(\sigma') = I$ and $J(\sigma') = J$. Thus, either $\sigma' = \sigma$ or $\sigma' = \sigma_{i_s,i_t}$.

(ii) Now assume that $s = 1$. Consider $\pi_{2,n}(\sigma)$, $\pi_{2,n}(\sigma')$ and $\pi_{2,n}(\sigma_{1,i_t})$. Note that $I(\pi_{2,n}(\sigma')) = I \setminus \{1\} = I(\pi_{2,n}(\sigma)) = I(\pi_{2,n}(\sigma_{1,i_t}))$.

   Further note that for any $m \neq t$ we get by the definition of $M_{[1,j_s]}(\sigma)$ that either $j_m < j_t$, or $i_m > i_t$ thus by $(***)$ $(R_{\sigma_{1,i_t}})_{p,r} = (R_\sigma)_{p,r}$ for $p = i_m, i_m + 1$ and $r = j_m, j_m - 1$. This forces also $(R_{\sigma'})_{p,r} = (R_\sigma)_{p,r}$ for $p = i_m, i_m + 1$ and $r = j_m, j_m - 1$. By Lemma 3.1 $j'_m = j_m$ for any $m \ : \ m > 1$ and $m \neq t$. In other words for any $m \neq 1, t$ one has $(i_m, j_m) \in \sigma'$. Since $I(\sigma') = I$ and $J(\sigma') = J$ we get that either $(1, j_1), (i_t, j_t) \in \sigma'$ (that is $\sigma' = \sigma$) or $(1, j_t), (i_t, j_1) \in \sigma'$ (that is $\sigma' = \sigma_{1,i_t}$).

■

3.15.

Proposition. *For* $\sigma = (i_1, j_1) \dots (i_k, j_k)$ *in* $\mathbf{S}_n^2$

$$D_1(\sigma) = \coprod_{s=1}^{k} \sigma_{i_s \uparrow} \cup \coprod_{s=1}^{k} \sigma_{j_s \rightarrow} \cup \coprod_{s=2}^{k} C_{i_s \uparrow \rightarrow}(\sigma) \cup \coprod_{s=1}^{k} C_{i_s \uparrow \downarrow}(\sigma)$$

Proof.

Let us denote $D'(\sigma) := \coprod\limits_{s=1}^{k} \sigma_{i_s \uparrow} \cup \coprod\limits_{s=1}^{k} \sigma_{j_s \rightarrow}$ and $D''(\sigma) := \coprod\limits_{s=2}^{k} C_{i_s \uparrow \rightarrow}(\sigma) \cup \coprod\limits_{s=1}^{k} C_{i_s \uparrow \downarrow}(\sigma)$. By lemmas 3.11-3.14 it remains to show that for any $\sigma' = (i'_1, j'_1) \dots (i'_k, j'_k)$ such that $\sigma' < \sigma$ there exist $\sigma'' \in D'(\sigma) \cup D''(\sigma)$ such that $\sigma'' \geq \sigma'$. This is a straightforward but rather long computation. It involves the comparison of $I(\sigma)$ and $J(\sigma)$ with $I(\sigma')$ and $J(\sigma')$. As in the proof of proposition 3.9 the argument is broken into the six claims.

Since $l(\sigma') = l(\sigma)$ one has that $I(\sigma') \cup J(\sigma') \neq I(\sigma) \cup J(\sigma)$ implies the existence of $q \in I(\sigma') \cup J(\sigma')$ such that $q \notin I(\sigma) \cup J(\sigma)$.

Claim 1. *Assume that there exists* $q \in I(\sigma')$ *such that* $q \notin I(\sigma) \cup J(\sigma)$. *Then there exists* $s \in M(\pi_{q+1,n}(\sigma))$ *such that* $\sigma_{i_s \uparrow} \geq \sigma'$.

Proof.

Note that $(R_{\sigma'})_{q,n} = (R_{\sigma'})_{q+1,n} + 1$ and $(R_\sigma)_{q,n} = (R_\sigma)_{q+1,n}$. Since $(R_{\sigma'})_{q,n} \leq (R_\sigma)_{q,n}$ this implies $(R_{\sigma'})_{q+1,n} < (R_\sigma)_{q+1,n}$ so that by Proposition 3.9 there exists $s \in M(\pi_{q+1,n}(\sigma))$ such that $(\pi_{q+1,n}(\sigma))^-_s \geq \pi_{q+1,n}(\sigma')$. Consider $\sigma_{i_s \uparrow}$. Since $s \in M(\pi_{q+1,n}(\sigma))$ and $q \notin I(\sigma)$ one has $\sigma_{i_s \uparrow} \neq \emptyset$. Moreover $m \geq q$. Hence,

$$(R_{\sigma_{i_s \uparrow}})_{i,j} \begin{cases} \geq (R_{(\pi_{q+1,n}(\sigma))^-_s})_{i,j}, & \text{if } i \geq q+1; \\ = (R_\sigma)_{i,j}, & \text{if } i \leq q. \end{cases}$$

Thus, $\sigma_{i_s \uparrow} \geq \sigma'$.

■

By symmetry of the rank matrix around the anti-diagonal we get the same result for the case $q \in J(\sigma')$. We need the complete formulation of this result in what follows.

Claim 2. *Assume that there exists* $q \in J(\sigma')$ *such that* $q \notin I(\sigma) \cup J(\sigma)$. *Then there exists* $r \in M(\pi_{1,q-1}(\sigma))$ *such that* $\sigma_{j_r \rightarrow} \geq \sigma'$.

Now we have to consider the case $I(\sigma') \cup J(\sigma') = I(\sigma) \cup J(\sigma)$. (Note that this does not imply $I(\omega) \cup J(\omega) = I(\sigma) \cup J(\sigma)$ for all $\omega \ : \ \sigma \geq \omega \geq \sigma''$). We show that in this case there exists $\sigma'' \in D'(\sigma) \cup D''(\sigma)$ such that $\sigma'' \geq \sigma'$. The proof is by induction. For $n = 4$ it holds trivially, as we have seen in 3.13, 3.14. Assume it for $n - 1$.

Claim 3. *Assume that* $I(\sigma') \cup J(\sigma') = I(\sigma) \cup J(\sigma)$. *Let* $n \notin J(\sigma)$. *Then there exists* $\sigma'' \in D'(\sigma) \cup D''(\sigma)$ *such that* $\sigma'' \geq \sigma'$.

Proof.

Since $n \notin J(\sigma), J(\sigma')$ one has $\pi_{1,n-1}(\sigma)=\sigma$ and $\pi_{1,n-1}(\sigma')=\sigma'$. Thus, by the induction hypothesis there exists $\sigma'' \in D'(\pi_{1,n-1}(\sigma)) \cup D''(\pi_{1,n-1}(\sigma)$ such that $\sigma'' \geq \pi_{1,n-1}(\sigma')$. Considering σ'' as an element of $\mathbf{S}_n^2$ we get the result.

■

Now assume that $n \in J(\sigma)$. Possibly changing the order of pairs we write $\sigma = (i_1, j_1)\dots(i_{k-1}, j_{k-1})(i, n)$ and $\sigma' = (i'_1, j'_1)\dots(i'_{k-1}, j'_{k-1})(i', n)$. Let us start with the case $i' \neq i$. In that case $i \in I(\pi_{1,n-1}(\sigma')) \cup J(\pi_{1,n-1}(\sigma'))$ so that either $i \in I(\pi_{1,n-1}(\sigma'))$ or $i \in J(\pi_{1,n-1}(\sigma'))$. We consider each case in a separate claim.

Claim 4. *Assume $I(\sigma') \cup J(\sigma') = I(\sigma) \cup J(\sigma)$. Let $(i,n) \in \sigma$, and $(i,n) \notin \sigma'$. Assume $i \in I(\sigma')$. Then there exists $i_t \in I(\sigma)$ such that $i_t > i$ and $\sigma_{i,i_t} \geq \sigma'$.*

Proof.

Since $i \notin I(\pi_{1,n-1}(\sigma))$ one has $(R_\sigma)_{i,n-1} = (R_\sigma)_{i+1,n-1}$. Since $i \in I(\sigma')$ and $(i,n) \notin \sigma'$ one has $i \in I(\pi_{1,n-1}(\sigma'))$. Thus, $(R_{\sigma'})_{i,n-1} = (R_{\sigma'})_{i+1,n-1} + 1$. Since $\sigma' < \sigma$, in particular, one has $(R_{\sigma'})_{i,n-1} \leq (R_\sigma)_{i,n-1}$. Therefore, $(R_{\sigma'})_{i+1,n-1} < (R_\sigma)_{i+1,n-1}$. By Proposition 3.9 there exists $t \in M(\pi_{i+1,n-1}(\sigma))$ such that $(\pi_{i+1,n-1}(\sigma))_t^- \geq \pi_{i+1,n-1}(\sigma')$. Note that $(\pi_{i+1,n-1}(\sigma))_t^- = \pi_{i+1,n-1}(\sigma_t^-)$. In particular, $(R_{\sigma'})_{p,q} \leq (R_{\sigma_t^-})_{p,q}$ whenever $p \geq i+1$ and $q \leq n-1$.

Note also that $t \in M_{[i,n]}(\sigma)$ so that σ_{i,i_t} exists. By 3.14 $(***')$

$$(R_{\sigma_{i,i_t}})_{p,q} = \begin{cases} (R_\sigma)_{p,q} & \text{if } p \leq i \text{ or } q = n; \\ (R_{\sigma_t^-})_{p,q} & \text{otherwise;} \end{cases}$$

Hence, $\sigma_{i,i_t} \geq \sigma'$.

■

Claim 5. *Assume $I(\sigma') \cup J(\sigma') = I(\sigma) \cup J(\sigma)$. Let $(i,n) \in \sigma$ and $i \in J(\sigma')$. Then there exists $r \in M(\pi_{1,i-1}(\sigma))$ such that $\sigma'' \geq \sigma'$ where $\sigma'' = \sigma_{j_r\to}$ or $\sigma'' = \sigma_{j_r,i}$.*

Proof.

Since $i \in J(\pi_{1,n-1}(\sigma'))$ and $i \notin I(\pi_{1,n-1}(\sigma)) \cup J(\pi_{1,n-1}(\sigma))$ by Claim 2 there exists $r \in M(\pi_{1,i-1}(\sigma))$ such that $(\pi_{1,n-1}(\sigma))_{j_r\to} \geq \pi_{1,n-1}(\sigma')$.

If there exists $r \in M(\pi_{1,i-1}(\sigma))$ such that $m < i$ then $\sigma_{j_r\to} \neq \emptyset$ and $\sigma_{j_r\to} \geq \sigma'$. Indeed, $\pi_{1,n-1}(\sigma_{j_r\to}) = (\pi_{1,n-1}(\sigma))_{j_r\to}$ so that $(R_{\sigma_{j_r\to}})_{p,q} \geq (R_{\sigma'})_{p,q}$ whenever $q < n$ and by $(*')$ of 3.12 $(R_{\sigma_{j_r\to}})_{p,n} = (R_\sigma)_{p,n} \geq (R_{\sigma'})_{p,n}$ for any p.

If for every $r \in M(\pi_{1,i-1}(\sigma))$ one has $m = i$ then $\sigma_{j_r,i}$ exists for every r. Let us take maximal such r. Let us show that $\sigma_{j_r,i} \geq \sigma'$.

(i) $(R_{\sigma_{j_r,i}})_{p,q} = (R_{(\pi_{1,n-1}(\sigma))_{j_r\to}})_{p,q} \geq (R_{\sigma'})_{p,q}$ whenever $q < n$.

(ii) By 3.13 $(**)$ $(R_{\sigma_{j_r,i}})_{p,n} = (R_\sigma)_{p,n}$ whenever either $p > i$ or $p \leq j_r$. Thus, $(R_{\sigma_{j_r,i}})_{p,n} \geq (R_{\sigma'})_{p,n}$ whenever either $p > i$ or $p \leq j_r$.
(iii) Since $i \in J(\sigma'), J(\sigma_{j_r,i})$ we get by the form of the rank matrix $(R_{\sigma'})_{i,n} = (R_{\sigma'})_{i+1,n}$ and $(R_{\sigma_{j_r,i}})_{i,n} = (R_{\sigma_{j_r,i}})_{i+1,n}$ so that $(R_{\sigma'})_{i,n} \leq (R_{\sigma_{j_r,i}})_{i,n}$.
(iv) Since r is maximal and $m = i$ by definition of $\sigma_{j_r\rightarrow}$ one has $t \in I(\sigma)$ for any $t \;:\; j_r < t < i$, thus, also $t \in I(\sigma_{j_r,i})$ for any such t. Therefore, for any $t \;:\; j_r < t < i$, one has $(R_{\sigma_{j_r,i}})_{t,n} = (R_{\sigma_{j_r,i}})_{i,n} + (t - i)$. On the other hand by the form of the rank matrix $(R_{\sigma'})_{t,n} \leq (R_{\sigma'})_{i,n} + (t - i)$, so that $(R_{\sigma'})_{t,n} \leq (R_{\sigma_{j_r,i}})_{t,n}$ for any $t \;:\; j_r < t < i$.

Together (i)-(iv) give us $\sigma_{j_r,i} \geq \sigma'$.

■

Claim 6. *Assume* $I(\sigma') \cup J(\sigma') = I(\sigma) \cup J(\sigma)$. *Let* $(i, n) \in \sigma, \sigma'$. *Then there exists* $\sigma'' \in D'(\sigma) \cup D''(\sigma)$ *such that* $\sigma'' \geq \sigma'$.

Proof.

Since $I(\pi_{1,n-1}(\sigma)) \cup J(\pi_{1,n-1}(\sigma)) = I(\pi_{1,n-1}(\sigma')) \cup J(\pi_{1,n-1}(\sigma'))$ by the induction hypothesis there exists $\hat\sigma \in D'(\pi_{1,n-1}(\sigma)) \cup D''(\pi_{1,n-1}(\sigma))$ such that $\hat\sigma \geq \pi_{1,n-1}(\sigma')$.

Assume $i \notin I(\hat\sigma) \cup J(\hat\sigma)$. Then $\hat\sigma(i, n) \in \mathbf{S}_n^2$ and $\sigma' \leq \hat\sigma(i, n) < \sigma$. Moreover,

(i) If $\hat\sigma = (\pi_{1,n-1}(\sigma))_{i_q,i_r}$ then either $i_r < i$ or $i_q > i$ so that $\hat\sigma(i, n) = \sigma_{i_q,i_r}$.
(ii) If $\hat\sigma = (\pi_{1,n-1}(\sigma))_{j_r,i_q}$ then either $i_q < i$ or $j_r > i$ so that $\hat\sigma(i, n) = \sigma_{j_r,i_q}$.
(iii) If $\hat\sigma = (\pi_{1,n-1}(\sigma))_{i_s\uparrow}$ and either $i_s < i$ or $m > i$ then $\hat\sigma(i, n) = \sigma_{i_s\uparrow}$.
(iv) If $\hat\sigma = (\pi_{1,n-1}(\sigma))_{j_s\rightarrow}$ and either $j_s > i$ or $m < i$ then $\hat\sigma(i, n) = \sigma_{i_s\rightarrow}$.

Setting $\sigma'' = \hat\sigma(i, n)$ we get the result.

It is left to consider the two cases when $i \in I(\hat\sigma) \cup J(\hat\sigma)$.

(i) If $\hat\sigma = (\pi_{1,n-1}(\sigma))_{i_s\uparrow}$ and $m = i$ let $\sigma'' = \sigma_{i,i_s}$. It exists since $s \in M_{[i,n]}(\sigma)$. Moreover, $\sigma_{i,i_s} \geq \sigma'$ since $(R_{\sigma_{i,i_s}})_{p,q} = (R_{\hat\sigma})_{p,q}$ whenever $q < n$ and $(R_{\sigma_{i,i_s}})_{p,n} = (R_\sigma)_{p,n}$ for any p.
(ii) If $\hat\sigma = (\pi_{1,n-1}(\sigma))_{j_s\rightarrow}$ and $m(j_s) = i$ we need a more subtle analysis. First note that in this case $(R_\sigma)_{1,i} > (R_{\sigma'})_{1,i}$. Indeed, $(R_{\hat\sigma})_{1,i} = (R_{\hat\sigma})_{1,i-1} + 1$ and $(R_{\sigma'})_{1,i} = (R_{\sigma'})_{1,i-1}$. Thus, $(R_{\hat\sigma})_{1,i-1} \geq (R_{\sigma'})_{1,i-1}$ provides $(R_\sigma)_{1,i} \geq (R_{\hat\sigma})_{1,i} > (R_{\sigma'})_{1,i}$. Since $(R_\sigma)_{1,n} = (R_{\sigma'})_{1,n}$ there exists $j > i$ such that
(a) $p \in J(\sigma')$ implies $p \in J(\sigma)$ for any $p \;:\; i < p < j$;
(b) $j \in J(\sigma')$ and $j \notin J(\sigma)$.
Since $I(\sigma) \cup J(\sigma) = I(\sigma') \cup J(\sigma')$ statements (a) and (b) are equivalent to
(a') $p \in I(\sigma)$ implies $p \in I(\sigma')$ for any $p \;:\; i < p < j$;
(b') $j \in I(\sigma)$ and $j \notin I(\sigma')$.
Thus by (b') $(R_\sigma)_{j,n-1} = (R_\sigma)_{j+1,n-1} + 1$ and $(R_{\sigma'})_{j,n-1} = (R_{\sigma'})_{j+1,n-1}$ so that $(R_\sigma)_{j,n-1} > (R_{\sigma'})_{j,n-1}$. By (a') this provides $(R_\sigma)_{i+1,n-1} > (R_{\sigma'})_{i+1,n-1}$.

Thus, by Proposition 3.9 there exists t such that $\pi_{i+1,n-1}(\sigma') \leq (\pi_{i+1,n-1}(\sigma))^-_t$ which implies by the definition that σ_{i,i_t} exists and $\sigma' \leq \sigma_{i,i_t}$.

■

This completes the proof of Proposition 3.15.

■

3.16. The two previous propositions involve in their proofs long and tricky combinatorics. At least I have not succeeded to find shorter and simpler proofs. On the other hand the proof of the following lemma is more straightforward.

Lemma. *For any $\sigma \in \mathbf{S}_n^2$ and for any $\sigma' \in D_1(\sigma) \cup D_2(\sigma)$ one has $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$.*

Proof.

Consider $\sigma = (i_1, j_1) \dots (i_k, j_k)$ and $\sigma' \in D_1(\sigma) \cup D_2(\sigma)$. Since the metric topology has more open sets than the Zariski topology it is enough to show that $N_{\sigma'} \in \overline{\mathcal{B}}_\sigma$ for the metric topology on $\operatorname{End} \mathbb{C}^n$. We construct a one parameter family of matrices $\{A_k\} \in \mathbf{B}$ such that $\lim\limits_{k\to\infty} A_k N_\sigma A_k^{-1} = N_{\sigma'}$ in the metric topology. Each A_k will be an elementary upper-triangular matrix or a product of elementary upper-triangular matrices. We construct them explicitly. Let $E_i(a)$ and $E_{i+j}(a)$ be the invertible matrices defined as follows

$$(E_i(a))_{s,t} = \begin{cases} a, & \text{if } s = t = i; \\ \delta_{s,t} & \text{otherwise.} \end{cases} \quad (E_{i+j}(a))_{s,t} = \begin{cases} a, & \text{if } s = i,\ t = j; \\ \delta_{s,t} & \text{otherwise.} \end{cases}$$

One checks that $N_{\sigma'} = \lim\limits_{k\to\infty} A_k N_\sigma A_k^{-1}$ where A_k are given as follows

(i) For $\sigma' = \sigma_s^-$ take $A_k = E_{i_s}(\frac{1}{k})$;
(ii) For $\sigma' = \sigma_{i_s}$ take $A_k = E_{i_s}(\frac{1}{k}) E_{m_s+i_s}(1)$;
(iii) For $\sigma' = \sigma_{j_s}$ take $A_k = E_{i_s}(\frac{1}{k}) E_{j_s+m_{j_s}}(-k)$;
(iv) For $\sigma' = \sigma_{j_r,i_s}$ take $A_k = E_{i_r}(-\frac{1}{k}) E_{i_s}(\frac{1}{k}) E_{j_r+i_s}(1)$;
(v) For $\sigma' = \sigma_{i_s,i_t}$ take $A_k = E_{j_t}(k) E_{j_s}(-\frac{1}{k} E_{i_s+i_t}(k) E_{j_t+j_s}(\frac{1}{k})$.

■

3.17. As a corollary of Lemma 3.6, Propositions 3.9 and 3.15 and Lemma 3.16 we get

Corollary. (i) *Given $\sigma, \sigma' \in \mathbf{S}_n^2$ one has $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$ iff $\mathcal{V}_{\sigma'} \subset \mathcal{V}_\sigma$. In particular $\overline{\mathcal{B}}_\sigma = \coprod\limits_{\sigma' \leq \sigma} \mathcal{B}_{\sigma'}$.*
(ii) $\overline{\mathcal{B}}_\sigma = \mathcal{B}_\sigma \coprod \bigcup\limits_{\sigma' \in D_1(\sigma) \cup D_2(\sigma)} \overline{\mathcal{B}}_{\sigma'}$.

Proof.

First of all note that by 2.5 and $\mathbf{B}$ stability of $\mathcal{V}_\sigma$ each $\mathcal{V}_\sigma$ is a union of some of the $\mathcal{B}_{\sigma'}$.

(i) By Lemma 3.6 $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$ implies $\mathcal{V}_{\sigma'} \subset \mathcal{V}_\sigma$. On the other hand, if $\mathcal{V}_{\sigma'} \subset \mathcal{V}_\sigma$ then there exists a chain $\sigma = \eta_1 < \eta_2, \ldots < \eta_s = \sigma'$ such that $\eta_{i+1} \in D_1(\eta_i) \cup D_2(\eta_i)$ for every $1 \leq i \leq s-1$. Thus, by Lemma 3.16 one has $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_{\eta_{s-1}} \subset \cdots \subset \overline{\mathcal{B}}_\sigma$.

(ii) By Lemma 3.16 $\bigcup\limits_{\sigma' \in D_1(\sigma) \cup D_2(\sigma)} \overline{\mathcal{B}}_{\sigma'} \subset \overline{\mathcal{B}}_\sigma$. On the other hand, if $\overline{\mathcal{B}}_{\sigma'} \subsetneq \overline{\mathcal{B}}_\sigma$ then $\sigma' < \sigma$ just by Lemma 3.6. Thus, by Propositions 3.9 and 3.15 there exists $\sigma'' \in D_1(\sigma) \cup D_2(\sigma)$ such that $\sigma' \leq \sigma''$. Thus, by (i) $\mathcal{B}_{\sigma'} \subset \overline{\mathcal{B}}_{\sigma''}$. This concludes the proof.

■

3.18. Now we can prove theorem 3.5. Let us recall it

Theorem. *For any $\sigma \in \mathbf{S}_n^2$ one has*

$$\overline{\mathcal{B}}_\sigma = \mathcal{V}_\sigma = \coprod_{\sigma' \leq \sigma} \mathcal{B}_{\sigma'}.$$

Proof.

By 3.4 $\overline{\mathcal{B}}_\sigma \subset \mathcal{V}_\sigma$. Let us show that $\dim \mathcal{V}_\sigma = \dim \mathcal{B}_\sigma$ and that $\mathcal{B}_\sigma$ is the only component in $\mathcal{V}_\sigma$ of maximal dimension. Indeed, since $\mathcal{B}_\sigma \subset \mathcal{V}_\sigma$ we get that $\dim \mathcal{V}_\sigma \geq \dim \mathcal{B}_\sigma$. For any σ' such that $\dim \mathcal{B}_{\sigma'} \geq \dim \mathcal{B}_\sigma$ we have $\mathcal{B}_{\sigma'} \not\in \overline{\mathcal{B}}_\sigma$. Thus by 3.17(i) one has $\sigma' \not< \sigma$. That means that there exists i, j such that $(R_{\sigma'})_{i,j} > (R_\sigma)_{i,j}$, so that $\mathcal{B}_{\sigma'} \not\subset \mathcal{V}_\sigma$. Hence, $\dim \mathcal{V}_\sigma = \dim \mathcal{B}_\sigma$ and $\mathcal{B}_\sigma$ is the only component of this dimension in $\mathcal{V}_\sigma$. This together with 3.8 and 3.11 gives:

$$\mathcal{V}_\sigma = \mathcal{B}_\sigma \coprod \bigcup_{\sigma' \in D_1(\sigma) \cup D_2(\sigma)} \mathcal{V}_{\sigma'}.$$

The result follows by induction on order. Indeed, it is immediate that $\mathcal{V}_\sigma = \overline{\mathcal{B}}_\sigma = \mathcal{B}_\sigma \coprod \{0\}$ for $\sigma = (1, n)$ which is the minimal non-zero orbit. Now assume that this is true for all $\sigma' > \sigma$ then

$$\mathcal{V}_\sigma = \mathcal{B}_\sigma \coprod \bigcup_{\sigma' \in D_1(\sigma) \cup D_2(\sigma)} \mathcal{V}_{\sigma'} = \mathcal{B}_\sigma \coprod \bigcup_{\sigma' \in D_1(\sigma) \cup D_2(\sigma)} \overline{\mathcal{B}}_{\sigma'} = \overline{\mathcal{B}}_\sigma = \coprod \sigma' \geq \sigma \mathcal{B}_{\sigma'}.$$

■

4. **Results on orbital varieties of nilpotent order 2**

4.1. Let us apply the results of section 3 to the study of orbital varieties of nilpotent order 2 in $\mathfrak{sl}_n$. First we recall some facts about such orbital varieties and then in subsections 4.4, 4.5 formulate and prove the corollaries of section 3 for orbital varieties.

Recall the notion of a Young tableau from 1.5. We fill the boxes of $D_\lambda \in \mathbf{D}_n$ with n distinct positive integers (instead of integers $1, \ldots, n$) in such a manner that entries increase in rows from left to right and in columns from top to bottom. We call such an array a Young tableau or simply a tableau. If the numbers in a Young tableau form the set of integers from 1 to n, then the tableau is called a standard Young tableau. Again, for any Young tableau T associated to D_λ we put $\mathrm{sh}\,(T) = \lambda$.

Let $\mathbf{T}_n$ denote the set of all standard Young tableaux of size n.

Let $\mathbf{T}_n^2 \in \mathbf{T}_n$ be the subset of standard Young tableaux with two columns. Let $T = (T_1, T_2)$, where $T_1 = \begin{pmatrix} t_{1,1} \\ \vdots \\ t_{l,1} \end{pmatrix}$ is the first column of T and $T_2 = \begin{pmatrix} t_{1,2} \\ \vdots \\ t_{k,2} \end{pmatrix}$ is the second column of T.

Set $\sigma_{\scriptscriptstyle T} = (i_1, j_1) \dots (i_k, j_k)$ where $j_s = t_{s,2}$; $i_1 = t_{1,2} - 1$, and $i_s = \max\{d \in T_1 \setminus \{i_1, \dots, i_{s-1}\} \mid d < j_s\}$ for any $s > 1$. Note that now we order the cycles of σ so that the second entries increase. For example, take

$$T = \boxed{\begin{array}{cc} 1 & 4 \\ 2 & 5 \\ 3 & 7 \\ 6 & 8 \end{array}}\,,$$

Then $\sigma_{\scriptscriptstyle T} = (3,4)(2,5)(6,7)(1,8)$.

Let us denote $\mathcal{B}_T := \mathcal{B}_{\sigma_T}$. Recall the notion of $\mathcal{U}_T$ from 1.5. As we explained there if an orbital variety is associated to a nilpotent orbit of nilpotent order 2, then it contains a dense $\mathbf{B}-$orbit. Explicitly by [11, 4.13] one has

Proposition. *For $T \in \mathbf{T}_n^2$ one has $\overline{\mathcal{B}}_T = \overline{\mathcal{U}}_T$.*

4.2. Let us induce geometric and algebraic orders, defined in 2.11 from $\mathbf{S}_n^2$ to $\mathbf{T}_n^2$. For $T, S \in \mathbf{T}_n^2$ set $T \overset{\mathrm{G}}{\leq} S$ if $\sigma_{\scriptscriptstyle T} \overset{\mathrm{G}}{\leq} \sigma_{\scriptscriptstyle S}$ and set $T \overset{\mathrm{A}}{\leq} S$ if $\sigma_{\scriptscriptstyle T} \overset{\mathrm{A}}{\leq} \sigma_{\scriptscriptstyle S}$. As a straightforward corollary of [14, 3.7] we get that these two orders coincide, that is

Theorem. *For $T, S \in \mathbf{T}_n^2$ one has $T \overset{\mathrm{G}}{\leq} S$ iff $T \overset{\mathrm{A}}{\leq} S$.*

Since these two orders coincide on $\mathbf{T}_n^2$ we can define a partial order on $\mathbf{T}_n^2$ by taking $T \leq S$ if $T \overset{\mathrm{G}}{\leq} S$.

4.3. For $T \in \mathbf{T}_n^2$ set $\mathcal{D}(T)$ to be what is called in combinatorics the cover of T, that is

$$\mathcal{D}(T) := \{S \mid S > T \text{ and if } S \geq U > T \text{ then } U = S\}.$$

In [14, 3.18] $\mathcal{D}(T)$ is described explicitly.

Given $T = (T_1, T_2)$ where $T_1 = \begin{pmatrix} t_{1,1} \\ \vdots \\ t_{l,1} \end{pmatrix}$ is the first column of T and $T_2 = \begin{pmatrix} t_{1,2} \\ \vdots \\ t_{k,2} \end{pmatrix}$ is the second column of T. For $t_{s,2}$ let $j_s := \max\{j \mid t_{j,1} < t_{s,2}\}$. Set $T\langle t_{s,2}\rangle$ to be the tableau obtained from T by moving $t_{s,2}$ from the second column to the first one, that is $T\langle t_{s,2}\rangle := (T_1', T_2')$, where

$$T_1' = \begin{pmatrix} t_{1,1} \\ \vdots \\ t_{j_s,1} \\ t_{s,2} \\ t_{j_s+1,1} \\ \vdots \\ t_{l,1} \end{pmatrix} \qquad \text{and} \qquad T_2' = \begin{pmatrix} t_{1,2} \\ \vdots \\ t_{s-1,2} \\ t_{s+1,2} \\ \vdots \\ t_{k,2} \end{pmatrix}.$$

For example, take T from 4.1. Then

$$T\langle 4\rangle = \begin{array}{|cc|} \hline 1 & 5 \\ 2 & 7 \\ 3 & 8 \\ 4 & \\ 6 & \\ \hline \end{array}\,, \quad T\langle 5\rangle = \begin{array}{|cc|} \hline 1 & 4 \\ 2 & 7 \\ 3 & 8 \\ 5 & \\ 6 & \\ \hline \end{array}\,, \quad T\langle 7\rangle = \begin{array}{|cc|} \hline 1 & 4 \\ 2 & 5 \\ 3 & 8 \\ 6 & \\ 7 & \\ \hline \end{array}\,, \quad T\langle 8\rangle = \begin{array}{|cc|} \hline 1 & 4 \\ 2 & 5 \\ 3 & 7 \\ 6 & \\ 8 & \\ \hline \end{array}\,.$$

Note that $T\langle t_{s,2}\rangle$ is always a Young tableau.

Recall σ_s^- from 3.8. Let $\sigma_T = (i_1, t_{1,2}) \dots (i_k, t_{k,2})$. Let

$$\sigma_{T\langle t_{s,2}\rangle} = (i_1', t_{1,2}) \dots (i_{s-1}', t_{s,2})(i_{s+1}', t_{s+1,2}) \dots (i_k', t_{k,2}).$$

Note that for any $p < s$ one has $i_p' = i_p$. For $p > s$ one has $i_p' = i_p$ if and only if $i_p' > t_{s,2}$. In other words, $\sigma_{T\langle t_{s,2}\rangle} = (\sigma_T)_s^-$ if and only if for any $p > s$ one has $i_p > t_{s,2}$. This note permits us to reformulate [14, 3.18] as follows

Theorem. *For any* $T \in \mathbf{T}_n^2$ *one has*

$$\mathcal{D}(T) = \{T\langle t_{s,2}\rangle \mid \sigma_{T\langle t_{s,2}\rangle} = (\sigma_T)_s^-\}$$

For example, for T from 4.1 one has

$$\mathcal{D}(T)=\{T\langle 8\rangle\}=\left\{\begin{array}{|c|c|}\hline 1 & 4\\ 2 & 5\\ 3 & 7\\ \cline{2-2} 6 & \multicolumn{1}{c}{}\\ 8 & \multicolumn{1}{c}{}\\ \cline{1-1}\end{array}\right\}$$

4.4. As a straightforward corollary of Proposition 4.1 and Theorem 3.5 we get

Corollary. *For any* $T\in\mathbf{T}_n^2$ *the ideal of definition* $I(\mathcal{U}_T)$ *is the radical of* $\mathcal{I}_{\mathcal{B}_T}$.

$$I(\mathcal{U}_T)=\sqrt{\mathcal{I}_{\mathcal{B}_T}}.$$

Proof.

Indeed, by 4.1 $\overline{\mathcal{U}}_T=\overline{\mathcal{B}}_T$ and by 3.5 $\mathcal{V}(\mathcal{I}_{\sigma_T})=\overline{\mathcal{B}}_T$ thus $I(\mathcal{U}_T)=\sqrt{\mathcal{I}_{\mathcal{B}_T}}$.

■

4.5. We get also

Theorem. *For* $T\in\mathbf{T}_n^2$ *one has*

$$\overline{\mathcal{U}}_T=\bigcup_{S\geq T}\mathcal{U}_S$$

Proof.

Since $\mathcal{B}_T$ is dense in $\mathcal{U}_T$ we get

$$\mathcal{U}_T=\overline{\mathcal{B}}_T\cap\mathcal{O}_T=\coprod_{\substack{\sigma'\leq\sigma\\ l(\sigma')=l(\sigma)}}\mathcal{B}_{\sigma'}.$$

On the other hand by theorem 3.5

$$\overline{\mathcal{B}}_T=\coprod_{\sigma'\leq\sigma}\mathcal{B}_{\sigma'}=\coprod_{\substack{\sigma'\leq\sigma\\ l(\sigma')=l(\sigma)}}\mathcal{B}_{\sigma'}\coprod\coprod_{\substack{\sigma'<\sigma\\ l(\sigma')<l(\sigma)}}\mathcal{B}_{\sigma'}.$$

Further, by definition of $D_2(\sigma)$ we get that for any σ' such that $\sigma'<\sigma$ and $l(\sigma')<l(\sigma)$ there exists $\sigma''\in D_2(\sigma)$ such that $\sigma''\geq\sigma'$. In other words, by theorem 3.5, $\mathcal{B}_{\sigma'}\subset\overline{\mathcal{B}}_{\sigma''}$. Thus, we get

$$\overline{\mathcal{U}}_T=\overline{\mathcal{B}}_T=\mathcal{U}_T\coprod\bigcup_{\sigma'\in D_2(\sigma_T)}\overline{\mathcal{B}}_{\sigma'}.$$

Let us show that $D_2(\sigma_T)=\{\sigma_S\}_{S\in\mathcal{D}(T)}$. Indeed, take $T=(T_1,T_2)$ where $T_1=\begin{pmatrix} t_{1,1}\\ \vdots \\ t_{l,1}\end{pmatrix}$ and $T_2=\begin{pmatrix} t_{1,2}\\ \vdots \\ t_{k,2}\end{pmatrix}$. Then $\sigma_T=(t_{i_1,1},t_{1,2})\ldots(t_{i_k,1},t_{k,2})$ where $t_{i_1,1}=\max\{t_{s,1}\ :\ t_{s,1}<a_{1,2}\}$ and $t_{i_j,1}=\max\{t_{s,1}\ :\ t_{s,1}\in T_1\setminus\{t_{i_1,1},\ldots,t_{i_{j-1},1}\},\ t_{s,1}<t_{j,k}\}$ whenever $j>1$. Since now the cycles of σ_T are ordered so that the second entries increase, one has $j\in M(\sigma_T)$ $\iff t_{i_j,1}=\max\{t_{i_1,1},\ldots,t_{i_j,1}\}$ so that $j\in M(\sigma_T)$ if and only if $(\sigma_T)^-_j=\sigma_{T\langle t_{j,2}\rangle}$ for some $T\langle t_{j,2}\rangle\in\mathcal{D}(T)$. Thus, we get $\overline{\mathcal{B}}_T=\mathcal{U}_T\coprod\bigcup\limits_{S\in\mathcal{D}(T)}\overline{\mathcal{U}}_S$.

The proof is completed by the induction on rank of $\mathcal{U}_T$. If $\lambda(T)=(n-1,1)$ that is Rank$\mathcal{U}_T=1$ it is obvious that $\overline{\mathcal{U}}_T=\mathcal{U}_T\coprod\{0\}=\bigcup\limits_{S\geq T}\mathcal{U}_S$. Assume that this is true for S of rank less than k. Let T be of rank k. Note that that for any $S>T$ one has Rank$\,(S)\leq k-1$. Thus,

$$\overline{\mathcal{U}}_T=\mathcal{U}_T\coprod\bigcup_{S\in\mathcal{D}(T)}\overline{\mathcal{U}}_S=\mathcal{U}_T\coprod\bigcup_{S\in\mathcal{D}(T)}\left(\bigcup_{U\geq S}\mathcal{U}_U\right)=\bigcup_{S\geq T}\mathcal{U}_S.$$

■

INDEX OF NOTATION

1.1	$\mathbf{G}$, $\mathbf{B}$, $\mathbf{B}_n$, $\mathfrak{g}$, $\mathfrak{n}$, $\mathfrak{n}_n$, $\mathfrak{h}$, $\mathfrak{n}^-$, $\mathfrak{n}_n^-$, $S(*)$, $\mathcal{O}_u$, $\mathcal{B}_u$, $\mathcal{X}_2$, $\mathbf{S}_n$, $\mathbf{S}_n^2$, $\mathcal{I}_{\mathcal{B}}$, $\mathcal{V}(I)$
1.2	$\mathcal{U}$
1.4	$\mathcal{W} \overset{\mathrm{G}}{\leq} \mathcal{U}$
1.5	λ^*, λ, $\mathcal{O}_\lambda$, D_λ, T, $\mathrm{sh}\,(T)$, $D(u)$, $\pi_{1,i}$, $D_i(u)$, $\varphi(u)$, ν_T, $\mathcal{U}_T$
1.6	$\pi_{i,n}$, $D^i(u)$, $\vartheta(u)$, ν^T, $\pi_{i,j}(T)$, $D_{i,j}(T)$, ν'_T, $\mathcal{I}_T$
2.1	$P(n)$, $\mathbf{D}_n$
2.2	$\mathrm{sh}\,(u)$, $\mathrm{sh}\,(\mathcal{O}_u)$, $\mathcal{O}_{(l,k)}$
2.3	$e_{i,j}$, R, R^+, Π, $\alpha_{i,j}, \alpha_i, X_{i,j}$, $\mathbf{P}_{\alpha_i}$, $\mathbf{P}_{\langle i,j\rangle}$, $M_{\langle i,j\rangle}$, $L_{\langle i,j\rangle}$, $\mathbf{B}_{\langle i,j\rangle}$, $\mathfrak{m}_{\langle i,j\rangle}$, $\mathfrak{l}_{\langle i,j\rangle}$, $\mathfrak{n}_{\langle i,j\rangle}$, $\pi_{i,j}$, $D_{i,j}(u)$, D_u, $D_{\mathcal{B}_u}$
2.4	W, s_α, s_i, $S_{\langle i,j\rangle}$, $D_{\langle i,j\rangle}$, $\pi_{i,j} : \mathbf{S}_n \to \mathbf{S}_{\langle i,j\rangle}$
2.5	N_σ, $\mathcal{B}_\sigma$
2.6	$\mathcal{O}_\sigma$, $l(\sigma)$
2.8	$\lambda \leq \mu$
2.9	$x_{i,j}$
2.11	$\sigma' \overset{\mathrm{G}}{\leq} \sigma$, $\sigma' \overset{\mathrm{A}}{\leq} \sigma$
2.13	$(i,j) \in \sigma$, $I(\sigma)$, $J(\sigma)$, $S_{i,j}(\sigma)$, $\pi_{i,j}(\sigma)$
3.1	R_u, $R_{\mathcal{B}_u}$, R_σ
3.3	$\mathbf{R}_n^2$
3.4	I_2, $\mathcal{I}_{R_\sigma}$, $\mathcal{I}_\sigma$, $\mathcal{V}_{R_\sigma}$, $\mathcal{V}_\sigma$
3.5	$R \leq R'$, $\sigma \leq \sigma'$
3.7	$D_1(\sigma)$, $D_2(\sigma)$, $M(\sigma)$,
3.8	σ_s^-
3.11	$\sigma_{i_s\uparrow}$
3.12	$\sigma_{j_s\rightarrow}$
3.13	σ_{j_r,i_s}, $C_{i_s\uparrow\rightarrow}(\sigma)$
3.14	σ_{i_s,i_t}, $C_{i_s\uparrow\downarrow}$
4.1	$\mathbf{T}_n$, $\mathbf{T}_n^2$, T_1, T_2, σ_T, $\mathcal{B}_T$
4.2	$T \leq S$
4.3	$\mathcal{D}(T)$, $T\langle i\rangle$

References

[1] W.Borho, J.-L. Brylinski, Differential operators on homogeneous spaces, Invent. math. 80 (1985), 1-68.

[2] E. Benlolo, Etude sur les variétés orbitales dans $\mathfrak{sl}_n(\mathbb{C})$, Thesis, Haifa University, 1990.

[3] A. Braverman, A. Joseph, The minimal Realization from De formation Theory, J. algebra 205 (1998), 13-36.

[4] D. Eisenbud, C. Huneke, W. Vasconcelos, Direct methods for primary decompositions, Invent. Math. 110 no 2, 1992, pp. 207-235.

[5] W. Hesselink, Singularities in the nilpotent scheme of a classical group, Trans. Am. Math. Soc. 222 (1976), 1-32.

[6] A. Joseph, On the variety of a highest weight module, J. of Algebra 88 (1984), 238-278.

[7] A. Joseph, Orbital varieties, Goldie rank polynomials and unitary highest weight modules, in B. Orsted and H. Schlichtkrull (eds.), Algebraic and Analytic Methods in Representation Theory, Perspectives in Mathematics, Vol.17, Academic Press, London, 1997.

[8] A. Joseph, Orbital varieties of the minimal orbit, Ann. scient. Ec. Norm. Sup, 4 serie, vol. 31, 1998, pp 17-45.

[9] A. Joseph, A. Melnikov, Quantization of hypersurface orbital varieties in $\mathfrak{sl}_n$, The orbit method in geometry and physics. In honor of A.A. Kirillov, series "Progress in Mathematics", 213, Birkhauser, 2003, 165-196.

[10] A. Melnikov, Irreducibility of the associated varieties of simple highest weight modules in $\mathfrak{sl}(n)$, C.R.A.S. I, 316 (1993), 53-57.

[11] A.Melnikov, Orbital varieties in $\mathfrak{sl}_n$ and the Smith conjecture, J. of Algebra, 200, 1998, pp. 1 - 31.

[12] A.Melnikov, B-orbits in solutions to the equation $X^2 = 2$ in triangular matrices, J. of Algebra, 223, 2000, pp. 101 - 108.

[13] A. Melnikov, On orbital variety closures in $\mathfrak{sl}_n$ II. Descendants of a Richardson orbital variety, J. of Algebra 271, 2004, pp 698-724.

[14] A.Melnikov, The combinatorics of orbital variety closures of nilpotent order 2 in $\mathfrak{sl}_n$, Electronic Journal of Combinatorics, vol. 12(1), 2005, R21.

[15] A. Melnikov, On varieties in an orbital variety closure in semisimple Lie algebras, ArXiv, math.RT/0409445

[16] M. P. Schützenberger, La correspondance de Robinson, LN in Math. **597** (1976), 59-113.

[17] N.Spaltenstein, The fixed point set of a unipotent transformation on the flag manifold, Proc. Konin. Nederl. Akad. 79 (1976), 452-456.

[18] N. Spaltenstein, Classes unipotentes de sous-groupes de Borel, LN in Math. 964 Springer-Verlag, 1982.

[19] R. Steinberg, On the desingularization of the unipotent variety, Invent. Math. 36 (1976), 209-224.

[20] R. Steinberg, An occurrence of the Robinson-Schensted correspondence, J. of Algebra 113 (1988), 523-528.

[21] M. A. van Leeuwen, The Robinson-Schensted and Schützenberger algorithms, Part II: Geometric interpretations, CWI report AM-R9209 (1992).

Department of Mathematics, University of Haifa, Haifa 31905, Israel

E-mail address: `melnikov@math.haifa.ac.il`